

Cofree Com-PreLie algebras

Loïc Foissy

*Fédération de Recherche Mathématique du Nord Pas de Calais FR 2956
Laboratoire de Mathématiques Pures et Appliquées Joseph Liouville
Université du Littoral Côte d'Opale-Centre Universitaire de la Mi-Voix
50, rue Ferdinand Buisson, CS 80699, 62228 Calais Cedex, France*

email: foissy@univ-littoral.fr

Abstract

A Com-PreLie bialgebra is a commutative bialgebra with an extra preLie product satisfying some compatibilities with the product and the coproduct. We here give examples of cofree Com-PreLie bialgebras, including all the ones such that the preLie product is homogeneous of degree ≥ -1 . We also give a graphical description of free unitary Com-PreLie algebras, explicit their canonical bialgebra structure and exhibit with the help of a rigidity theorem certain cofree quotients, including the Connes-Kreimer Hopf algebra of rooted trees. We finally prove that the dual of these bialgebras are also enveloping algebras of preLie algebras, combinatorially described.

AMS classification. 17D25 16T05 05C05

Contents

1	Reminders on Com-PreLie algebras	3
1.1	Definitions	3
1.2	Linear endomorphism on primitive elements	4
1.3	Extension of the pre-Lie product	5
2	Examples on shuffle algebras	6
2.1	Preliminary lemmas	6
2.2	PreLie products of positive degree	9
2.3	PreLie products of degree -1	10
3	Free Com-PreLie algebras and quotients	11
3.1	Description of free Com-PreLie algebras	11
3.2	Quotients of $UCP(\mathcal{D})$	14
3.3	PreLie structure of $UCP(\mathcal{D})$ and $CP(\mathcal{D})$	16
4	Bialgebra structures on free Com-PreLie algebras	19
4.1	Tensor product of Com-PreLie algebras	19
4.2	Coproduct on $UCP(\mathcal{D})$	21
4.3	An application: Connes-Moscovici subalgebras	24
4.4	A rigidity theorem for Com-PreLie bialgebras	26
4.5	Dual of $UCP(\mathcal{D})$ and $CP(\mathcal{D})$	29
4.6	Extension of the preLie product \diamond to all partitioned trees	32

Introduction

Com-PreLie bialgebras, introduced in [5, 6], are commutative bialgebras with an extra preLie product, compatible with the product and coproduct: see Definition 1 below. They appeared in Control Theory, as the Lie algebra of the group of Fliess operators [8] naturally owns a Com-PreLie bialgebra structure, and its underlying bialgebra is a shuffle Hopf algebra. Free (non unitary) Com-PreLie bialgebras were also described, in terms of partitioned rooted trees.

We here give examples of connected cofree Com-PreLie bialgebras. As cocommutative cofree bialgebras are, up to isomorphism, shuffle algebras $Sh(V) = (T(V), \sqcup, \Delta)$, where V is the space of primitive elements, we first characterize Com-PreLie bialgebras structures on $Sh(V)$ in term of operators $\varpi : T(V) \otimes T(V) \rightarrow V$, satisfying two identities, see Proposition 8. In particular, if we assume that the obtained preLie bracket is homogeneous of degree 0 for the graduation of $Sh(V)$ by the length, then ϖ is reduced to a linear map $f : V \rightarrow V$, and the obtained preLie product is given by (Proposition 9):

$$\forall x_1, \dots, x_m, y_1, \dots, y_n \in V, \quad x_1 \dots x_m \bullet y_1 \dots y_n = \sum_{i=0}^n x_1 \dots x_{i-1} f(x_i) (x_{i+1} \dots x_m \sqcup y_1 \dots y_n).$$

In particular, if $V = Vect(x_0, x_1)$ and f is defined by $f(x_0) = 0$ and $f(x_1) = x_0$, we obtain the Com-PreLie bialgebra of Fliess operators in dimension 1. If we assume that the obtained preLie bracket is homogeneous of degree -1 , then ϖ is given by two bilinear products $*$ and $\{-, -\}$ on V such that $*$ is preLie, $\{-, -\}$ is antisymmetric and for all $x, y, z \in V$:

$$\begin{aligned} x * \{y, z\} &= \{x * y, z\}, \\ \{x, y\} * z &= \{x * y, z\} + \{x, y * z\} + \{\{x, y\}, z\}. \end{aligned}$$

This includes preLie products on V when $\{-, -\} = 0$ and nilpotent Lie algebras of nilpotency order 2 when $*$ = 0, see Proposition 11.

We then extend the construction of free Com-PreLie algebras of [5] in terms of partitioned trees (see Definition 12) to free unitary Com-PreLie algebras $UCP(\mathcal{D})$, with the help of a complementary decoration by integers. We obtain free Com-PreLie algebras $CP(\mathcal{D})$ as the augmentation ideal of a quotient of $UCP(\mathcal{D})$, the right action of the unit \emptyset on the generators of $UCP(\mathcal{D})$ being arbitrarily chosen (proposition 16). Recall that partitioned trees are rooted forests with an extra structure of a partition of its vertices into blocks; forgetting the blocks, we obtain the Connes-Kreimer Hopf algebra \mathcal{H}_{CK} of rooted trees [3, 4], which is given in this way a natural structure of Com-PreLie bialgebra (proposition 17). Using Livernet's rigidity theorem for preLie algebras, we prove that the augmentation ideals of $UCP(\mathcal{D})$ and $CP(\mathcal{D})$ are free as preLie algebras. Theorem 28 is a rigidity theorem which gives a simple criterion for a connected (as a coalgebra) Com-PreLie bialgebra to be cofree, in terms of the right action of the unit on its primitive elements. Applied to $CP(\mathcal{D})$ and \mathcal{H}_{CK} , it proves that they are isomorphic to shuffle bialgebras, which was already known for \mathcal{H}_{CK} . We also consider the dual Hopf algebras of $UCP(\mathcal{D})$ and $CP(\mathcal{D})$: as these Hopf algebras are right-sided combinatorial in the sense of [12], their duals are enveloping algebras of other preLie algebras, which we explicitly describe in Theorem 30, and then compare to the original Com-PreLie algebras.

This text is organized as follows: the first section contains reminders and lemmas on Com-PreLie algebras, including the extension of the Guin-Oudom extension of the preLie product in the Com-PreLie case. The second section deals with the characterization of preLie products on shuffle algebras. In the next section contains the description of free unitary Com-PreLie algebras and two families of quotients, whereas the fifth and last one contains results on the bialgebraic structures of these objects: existence of the coproduct, the rigidity theorem 28 and its applications, the dual preLie algebras, and an application to a family of subalgebras, named

Connes-Moscovici subalgebras.

Notations 1. 1. Let \mathbb{K} be a commutative field of characteristic zero. All the objects (vector spaces, algebras, coalgebras, PreLie algebras...) in this text will be taken over \mathbb{K} .

2. For all $n \in \mathbb{N}$, we denote by $[n]$ the set $\{1, \dots, n\}$. In particular, $[0] = \emptyset$.

1 Reminders on Com-PreLie algebras

Let V be a vector space.

- We denote by $T(V)$ the tensor algebra of V . Its unit is the empty word, which we denote by \emptyset . The element $v_1 \otimes \dots \otimes v_n \in V^{\otimes n}$, with $v_1, \dots, v_n \in V$, will be shortly denoted by $v_1 \dots v_n$. The deconcatenation coproduct of $T(V)$ is defined by:

$$\forall v_1, \dots, v_n \in V, \quad \Delta(v_1 \dots v_n) = \sum_{i=0}^n v_1 \dots v_i \otimes v_{i+1} \dots v_n.$$

The shuffle product of $T(V)$ is denoted by \sqcup . Recall that it can be inductively defined:

$$\forall x, y \in V, u, v \in T(V), \quad \emptyset \sqcup v = 0, \quad xu \sqcup yv = x(u \sqcup yv) + y(xu \sqcup v).$$

For example, if $v_1, v_2, v_3, v_4 \in V$:

$$\begin{aligned} v_1 \sqcup v_2 v_3 v_4 &= v_1 v_2 v_3 v_4 + v_2 v_1 v_3 v_4 + v_2 v_3 v_1 v_4 + v_2 v_3 v_4 v_1, \\ v_1 v_2 \sqcup v_3 v_4 &= v_1 v_2 v_3 v_4 + v_1 v_3 v_2 v_4 + v_1 v_3 v_4 v_2 + v_3 v_1 v_2 v_4 + v_3 v_1 v_4 v_2 + v_3 v_4 v_1 v_2, \\ v_1 v_2 v_3 \sqcup v_4 &= v_1 v_2 v_3 v_4 + v_1 v_2 v_4 v_3 + v_1 v_2 v_4 v_3 + v_1 v_4 v_2 v_3 + v_4 v_1 v_2 v_3. \end{aligned}$$

$Sh(V) = (T(V), \sqcup, \Delta)$ is a Hopf algebra, known as the shuffle algebra of V .

- $S(V)$ is the symmetric algebra of V . It is a Hopf algebra, with the coproduct defined by:

$$\forall v \in V, \quad \Delta(v) = v \otimes \emptyset + \emptyset \otimes v.$$

- $coS(V)$ is the subalgebra of $(T(V), \sqcup)$ generated by V . It is the greatest cocommutative Hopf subalgebra of $(T(V), \sqcup, \Delta)$, and is isomorphic to $S(V)$ via the following algebra morphism:

$$\theta : \begin{cases} (S(V), m, \Delta) & \longrightarrow & (coS(V), \sqcup, \Delta) \\ v_1 \dots v_k & \longrightarrow & v_1 \sqcup \dots \sqcup v_k. \end{cases}$$

1.1 Definitions

Definition 1. 1. A Com-PreLie algebra [5, 6] is a family $A = (A, \cdot, \bullet)$, where A is a vector space, \cdot and \bullet are bilinear products on A , such that:

$$\begin{aligned} \forall a, b \in A, & \quad a \cdot b = b \cdot a, \\ \forall a, b, c \in A, & \quad (a \cdot b) \cdot c = a \cdot (b \cdot c), \\ \forall a, b, c \in A, & \quad (a \bullet b) \bullet c - a \bullet (b \bullet c) = (a \bullet c) \bullet b - a \bullet (c \bullet b) \quad (\text{preLie identity}), \\ \forall a, b, c \in A, & \quad (a \cdot b) \bullet c = (a \bullet c) \cdot b + a \cdot (b \bullet c) \quad (\text{Leibniz identity}). \end{aligned}$$

In particular, (A, \cdot) is an associative, commutative algebra and (A, \bullet) is a right preLie algebra. We shall say that A is unitary if the algebra (A, \cdot) is unitary.

2. A Com-PreLie bialgebra is a family $(A, \cdot, \bullet, \Delta)$, such that:

(a) (A, \cdot, \bullet) is a Com-PreLie algebra.

(b) (A, \cdot, Δ) is a bialgebra.

(c) For all $a, b \in A$:

$$\Delta(a \bullet b) = a^{(1)} \otimes a^{(2)} \bullet b + a^{(1)} \bullet b^{(1)} \otimes a^{(2)} \cdot b^{(2)},$$

with Sweedler's notation $\Delta(x) = x^{(1)} \otimes x^{(2)}$.

Remark 1. If $(A, \cdot, \bullet, \Delta)$ is a Com-PreLie bialgebra, then for any $\lambda \in \mathbb{K}$, $(A, \cdot, \lambda \bullet, \Delta)$ also is.

Lemma 2. 1. Let (A, \cdot, \bullet) be a unitary Com-PreLie algebra. Its unit is denoted by \emptyset . For all $a \in A$, $\emptyset \bullet a = 0$.

2. Let A be a Com-PreLie bialgebra, with counit ε . For all $a, b \in A$, $\varepsilon(a \bullet b) = 0$.

Proof. 1. Indeed, $\emptyset \bullet a = (\emptyset \cdot \emptyset) \bullet a = (\emptyset \bullet a) \cdot \emptyset + \emptyset \cdot (\emptyset \bullet a) = 2(\emptyset \bullet a)$, so $\emptyset \bullet a = 0$.

2. For all $a, b \in A$:

$$\begin{aligned} \varepsilon(a \bullet b) &= (\varepsilon \otimes \varepsilon) \circ \Delta(a \bullet b) \\ &= \varepsilon(a^{(1)})\varepsilon(a^{(2)} \bullet b) + \varepsilon(a^{(1)} \bullet b^{(1)})\varepsilon(a^{(2)} \cdot b^{(2)}) \\ &= \varepsilon(a^{(1)})\varepsilon(a^{(2)} \bullet b) + \varepsilon(a^{(1)} \bullet b^{(1)})\varepsilon(a^{(2)})\varepsilon(b^{(2)}) \\ &= \varepsilon(a \bullet b) + \varepsilon(a \bullet b), \end{aligned}$$

so $\varepsilon(a \bullet b) = 0$. □

Remark 2. Consequently, if a is primitive:

$$\Delta(a \bullet b) = \emptyset \otimes a \bullet b + a \bullet b^{(1)} \otimes b^{(2)}.$$

The map $b \mapsto a \bullet b$ is a 1-cocycle for the Cartier-Quillen cohomology [3].

1.2 Linear endomorphism on primitive elements

If A is a Com-PreLie bialgebra, we denote by $Prim(A)$ the space of its primitive elements.

Proposition 3. Let A be a Com-PreLie bialgebra. Its unit is denoted by \emptyset .

1. If $x \in Prim(A)$, then $x \bullet \emptyset \in Prim(A)$. We denote by f_A the map:

$$f_A : \begin{cases} Prim(A) & \longrightarrow & Prim(A) \\ a & \longrightarrow & a \bullet \emptyset. \end{cases}$$

2. $Prim(A)$ is a preLie subalgebra of (A, \bullet) if, and only if, $f_A = 0$.

Proof. 1. Indeed, if a is primitive:

$$\Delta(a \bullet \emptyset) = a \otimes \emptyset \bullet \emptyset + \emptyset \otimes a \bullet \emptyset + a \bullet \emptyset \otimes \emptyset \cdot \emptyset + \emptyset \bullet \emptyset \otimes a \cdot \emptyset = 0 + \emptyset \otimes \emptyset \bullet a + a \bullet \emptyset \otimes \emptyset + 0,$$

so $a \bullet \emptyset$ is primitive.

2. and 3. Let $a, b \in Prim(A)$.

$$\begin{aligned} \Delta(a \bullet b) &= a \otimes \emptyset \bullet b + \emptyset \otimes a \bullet b + \emptyset \bullet \emptyset \otimes a \cdot b + a \bullet \emptyset \otimes b + \emptyset \bullet b \otimes a + a \bullet b \otimes \emptyset \\ &= \emptyset \otimes a \bullet b + a \bullet b \otimes \emptyset + f_A(a) \otimes b. \end{aligned}$$

Hence, $Prim(A)$ is a preLie subalgebra if, and only if, for any $a, b \in A$, $f_A(a) \otimes b = 0$, that is to say if, and only if, $f_A = 0$. □

1.3 Extension of the pre-Lie product

Let A be a Com-PreLie algebra. It is a Lie algebra, with the bracket defined by:

$$\forall x, y \in A, [x, y] = x \bullet y - y \bullet x.$$

We shall use the Oudom-Guin construction of its enveloping algebra [13, 14]. In order to avoid confusions, we shall denote by \times the usual product of $S(A)$ and by 1 its unit. We extend the preLie product \bullet into a product from $S(A) \otimes S(A)$ into $S(A)$:

1. If $a_1, \dots, a_k \in A$, $(a_1 \times \dots \times a_k) \bullet 1 = a_1 \times \dots \times a_k$.
2. If $a, a_1, \dots, a_k \in A$:

$$a \bullet (a_1 \times \dots \times a_k) = (a \bullet (a_1 \times \dots \times a_{k-1})) \bullet a_k - \sum_{i=1}^{k-1} a \bullet (a_1 \times \dots \times (a_i \bullet a_k) \times \dots \times a_{k-1}).$$

3. If $x, y, z \in S(A)$, $(x \times y) \bullet z = (x \bullet z^{(1)}) \times (y \bullet z^{(2)})$, where $\Delta(z) = z^{(1)} \otimes z^{(2)}$ is the usual coproduct of $S(A)$.

Notations 2. If $c_1, \dots, c_n \in A$ and $I = \{i_1, \dots, i_k\} \subseteq [n]$, we put:

$$\prod_{i \in I}^{\times} c_i = c_{i_1} \times \dots \times c_{i_k}.$$

Proposition 4. 1. Let A be a Com-PreLie algebra. If $a, b, c_1, \dots, c_n \in A$:

$$(a \cdot b) \bullet (c_1 \times \dots \times c_k) = \sum_{I \subseteq [n]} \left(a \bullet \prod_{i \in I}^{\times} c_i \right) \cdot \left(b \bullet \prod_{i \notin I}^{\times} c_i \right).$$

2. Let A be a Com-PreLie bialgebra. If $a, b_1, \dots, b_n \in A$:

$$\Delta(a \bullet (b_1 \times \dots \times b_n)) = \sum_{I \subseteq [n]} a^{(1)} \bullet \left(\prod_{i \in I}^{\times} b_i^{(1)} \right) \otimes \left(\prod_{i \in I}^{\times} b_i^{(2)} \right) a^{(2)} \bullet \left(\prod_{i \notin I}^{\times} b_i \right).$$

Proof. These are proved by direct, but quite long, inductions on n . □

Lemma 5. Let A be a Com-PreLie bialgebra. For all $a \in \text{Prim}(A)$, $k \geq 0$, $b_1, \dots, b_l \in A$:

$$a \bullet \emptyset^{\times k} \times b_1 \times \dots \times b_l = f_A^k(a) \bullet b_1 \times \dots \times b_l.$$

Proof. This is obvious if $k = 0$. Let us prove it for $k = 1$ by induction on l . It is obvious if $l = 0$. Let us assume the result at rank $l - 1$. Then:

$$\begin{aligned} a \bullet \emptyset \times b_1 \times \dots \times b_l &= (a \bullet \emptyset \times b_1 \times \dots \times b_{l-1}) \bullet b_l + a \bullet (\emptyset \bullet b_l) \times b_1 \times \dots \times b_{l-1} \\ &+ \sum_{i=1}^{l-1} a \bullet \emptyset \times b_1 \times \dots \times (b_i \bullet b_l) \times \dots \times b_{l-1} \\ &= (f_A(a) \bullet b_1 \times \dots \times b_{l-1}) \bullet b_l + 0 + \sum_{i=1}^{l-1} f_A(a) \bullet b_1 \times \dots \times (b_i \bullet b_l) \times \dots \times b_{l-1} \\ &= f_A(a) \bullet b_1 \times \dots \times b_l. \end{aligned}$$

The result is proved for $k \geq 2$ by an induction on k . □

2 Examples on shuffle algebras

2.1 Preliminary lemmas

We shall denote by $\pi : T(V) \rightarrow V$ the canonical projection.

Lemma 6. *Let $\varpi : T(V) \otimes T(V) \rightarrow V$ be a linear map.*

1. *There exists a unique map $\bullet : T(V) \otimes T(V) \rightarrow T(V)$ such that:*

(a) $\pi \circ \bullet = \varpi.$

(b) *For all $u, v \in T(V)$:*

$$\Delta(u \bullet v) = u^{(1)} \otimes u^{(2)} \bullet v + u^{(1)} \bullet v^{(1)} \otimes u^{(2)} \sqcup v^{(2)}. \quad (1)$$

This product \bullet is given by:

$$\forall u, v \in T(V), \quad u \bullet v = u^{(1)} \varpi(u^{(2)} \otimes v^{(1)})(u^{(3)} \sqcup v^{(2)}).$$

2. *The following conditions are equivalent:*

(a) *For all $u, v, w \in T(V)$:*

$$(u \sqcup v) \bullet w = (u \bullet w) \sqcup v + u \sqcup (v \bullet w).$$

(b) *For all $u, v, w \in T(V)$:*

$$\varpi((u \sqcup v) \otimes w) = \varepsilon(u) \varpi(v \otimes w) + \varepsilon(v) \varpi(u \otimes w). \quad (2)$$

3. *Let $N \in \mathbb{Z}$. The following conditions are equivalent:*

(a) *\bullet is homogeneous of degree N , that is to say:*

$$\forall k, l \geq 0, \quad V^{\otimes k} \bullet V^{\otimes l} \subseteq V^{\otimes(k+l+N)}.$$

(b) *For all $k, l \geq 0$, such that $k + l + N \neq 1$, $\varpi(V^{\otimes k} \otimes V^{\otimes l}) = (0)$.*

We use the convention $V^{\otimes p} = (0)$ if $p < 0$.

Proof. 1. *Existence.* Let \bullet be the product on $T(V)$ defined by:

$$\forall u, v \in T(V), \quad u \bullet v = u^{(1)} \varpi(u^{(2)} \otimes v^{(1)})(u^{(3)} \sqcup v^{(2)}).$$

As ϖ takes its values in V , for all $u, v \in T(V)$:

$$\begin{aligned} \pi(u \bullet v) &= \varepsilon(u^{(1)}) \varpi(u^{(2)} \otimes v^{(1)}) \varepsilon(u^{(3)} \sqcup v^{(2)}) \\ &= \varepsilon(u^{(1)}) \varpi(u^{(2)} \otimes v^{(1)}) \varepsilon(u^{(3)}) \varepsilon(v^{(2)}) \\ &= \varpi(u \otimes v). \end{aligned}$$

We denote by m the concatenation product of $T(V)$. As $(T(V), m, \Delta)$ is an infinitesimal bialgebra [10, 11], for all $u, v \in T(V)$:

$$\begin{aligned} \Delta(u \bullet v) &= u^{(1)} \otimes u^{(2)} \varpi(u^{(3)} \otimes v^{(1)})(u^{(4)} \sqcup v^{(2)}) + u^{(1)} \varpi(u^{(2)} \otimes v^{(1)}) \otimes u^{(3)} \sqcup v^{(2)} \\ &\quad + u^{(1)} \otimes \varpi(u^{(2)} \otimes v^{(1)})(u^{(3)} \sqcup v^{(2)}) + u^{(1)} \varpi(u^{(2)} \otimes v^{(1)})(u^{(3)} \sqcup v^{(2)}) \otimes u^{(4)} \sqcup v^{(3)} \\ &\quad - u^{(1)} \varpi(u^{(2)} \otimes v^{(1)}) \otimes u^{(3)} \sqcup v^{(2)} - u^{(1)} \otimes \varpi(u^{(2)} \otimes v^{(1)})(u^{(3)} \otimes v^{(2)}) \\ &= u^{(1)} \otimes u^{(2)} \varpi(u^{(3)} \otimes v^{(1)})(u^{(4)} \sqcup v^{(2)}) + u^{(1)} \varpi(u^{(2)} \otimes v^{(1)})(u^{(3)} \sqcup v^{(2)}) \otimes u^{(4)} \sqcup v^{(3)} \\ &= u^{(1)} \otimes u^{(2)} \bullet v + u^{(1)} \bullet v^{(1)} \otimes u^{(2)} \sqcup v^{(2)}. \end{aligned}$$

Unicity. Let \diamond be another product satisfying the required properties. Let us denote that $u \diamond v = u \bullet v$ for any words u, v of respective lengths k and l . If $k = 0$, then we can assume that $u = \emptyset$. We proceed by induction on l . If $l = 0$, then we can assume that $v = \emptyset$. By (1), $\emptyset \bullet \emptyset$ and $\emptyset \diamond \emptyset$ are primitive elements of $T(V)$, so belong to V . Hence:

$$\emptyset \bullet \emptyset = \pi(\emptyset \bullet \emptyset) = \varpi(\emptyset \otimes \emptyset) = \pi(\emptyset \diamond \emptyset) = \emptyset \diamond \emptyset.$$

If $l \geq 1$, then, by (1):

$$\begin{aligned} \Delta(\emptyset \bullet v) &= \emptyset \otimes \emptyset \bullet v + \emptyset \bullet v \otimes \emptyset + \emptyset \bullet \emptyset \otimes v + \emptyset \bullet v' \otimes v'', \\ \tilde{\Delta}(\emptyset \bullet v) &= \emptyset \bullet \emptyset \otimes v + \emptyset \bullet v' \otimes v''. \end{aligned}$$

The same computation for \diamond and the induction hypothesis on l , applied to (\emptyset, v') , imply that $\tilde{\Delta}(\emptyset \bullet v - \emptyset \diamond v) = 0$, so $\emptyset \bullet v - \emptyset \diamond v \in V$. Finally:

$$\emptyset \bullet v - \emptyset \diamond v = \pi(\emptyset \bullet v - \emptyset \diamond v) = \varpi(\emptyset \otimes v - \emptyset \otimes v) = 0.$$

If $k \geq 1$, we proceed by induction on l . If $l = 0$, we can assume that $v = \emptyset$; (1) implies that $\tilde{\Delta}(u \bullet \emptyset - u \diamond \emptyset) = 0$, so $u \bullet \emptyset - u \diamond \emptyset = 0$ and, applying π , finally $u \bullet \emptyset = u \diamond \emptyset$. If $l \geq 1$, by (1), the induction hypothesis on k applied to (u', v) and the induction hypothesis on l applied to (u, \emptyset) and (u, v') :

$$\begin{aligned} \tilde{\Delta}(u \bullet v) &= u' \otimes u'' \bullet v + u \bullet \emptyset \otimes v + u \bullet v' \otimes v'' \\ &= u' \otimes u'' \diamond v + u \diamond \emptyset \otimes v + u \diamond v' \otimes v'' = \tilde{\Delta}(u \diamond v). \end{aligned}$$

As before, $u \bullet v = u \diamond v$.

2. \implies . As ϖ takes its values in V , we have:

$$\begin{aligned} \varpi(u \sqcup v) \otimes w &= \varpi((u \bullet w) \sqcup v + u \sqcup (v \bullet w)) \\ &= \varepsilon(v) \varpi(u \otimes w) + \varepsilon(u) \varpi(v \otimes w). \end{aligned}$$

\Leftarrow . For all $u, v, w \in T(V)$:

$$\begin{aligned} (u \sqcup v) \bullet w &= (u^{(1)} \sqcup v^{(1)}) \varpi((u^{(2)} \sqcup v^{(2)}) \otimes w^{(1)}) (u^{(3)} \sqcup v^{(3)} \sqcup w^{(2)}) \\ &= \varepsilon(u^{(2)}) (u^{(1)} \sqcup v^{(1)}) \varpi(v^{(2)} \otimes w^{(1)}) (u^{(3)} \sqcup v^{(3)} \sqcup w^{(2)}) \\ &\quad + \varepsilon(v^{(2)}) (u^{(1)} \sqcup v^{(1)}) \varpi(u^{(2)} \otimes w^{(1)}) (u^{(3)} \sqcup v^{(3)} \sqcup w^{(2)}) \\ &= (u^{(1)} \sqcup v^{(1)}) \varpi(v^{(2)} \otimes w^{(1)}) (u^{(2)} \sqcup v^{(3)} \sqcup w^{(2)}) \\ &\quad + (u^{(1)} \sqcup v^{(1)}) \varpi(u^{(2)} \otimes w^{(1)}) (u^{(3)} \sqcup v^{(2)} \sqcup w^{(2)}) \\ &= u \sqcup \left(v^{(1)} \varpi(v^{(2)} \otimes w^{(1)}) (v^{(3)} \sqcup w^{(2)}) \right) \\ &\quad + v \sqcup \left(u^{(1)} \varpi(u^{(2)} \otimes w^{(1)}) (u^{(3)} \sqcup w^{(2)}) \right) \\ &= u \sqcup (v \bullet w) + (u \bullet w) \sqcup v. \end{aligned}$$

So the compatibility between \sqcup and \bullet is satisfied.

3. Immediate. □

Remark 3. If (2) is satisfied, for $u = v = \emptyset$, we obtain:

$$\forall w \in T(V), \quad \varpi(\emptyset \otimes w) = 0.$$

Lemma 7. Let $\varpi : T(V) \otimes T(V) \longrightarrow V$, satisfying (2), and let \bullet be the product associated to ϖ in Lemma 6. Then $(T(V), \sqcup, \bullet, \Delta)$ is a Com-PreLie bialgebra if, and only if:

$$\forall u, v, w \in T(V), \quad \varpi(u \bullet v \otimes w) - \varpi(u \otimes v \bullet w) = \varpi(u \bullet w \otimes v) - \varpi(u \otimes w \bullet v). \quad (3)$$

Proof. \implies . This is immediately obtained by applying π to the preLie identity, as $\varpi = \pi \circ \bullet$.

\impliedby . By lemma 6, it remains to prove that \bullet is preLie. For any $u, v, w \in T(V)$, we put:

$$PL(u, v, w) = (u \bullet v) \bullet w - u \bullet (v \bullet w) - (u \bullet w) \bullet v + u \bullet (w \bullet v).$$

By hypothesis, $\pi \circ PL(u, v, w) = 0$ for any $u, v, w \in T(V)$. Let us prove that $PL(u, v, w) = 0$ for any $u, v, w \in T(V)$. A direct computation using (1) shows that:

$$\Delta(PL(u, v, w)) = u^{(1)} \otimes PL(u^{(2)}, v, w) \otimes u^{(1)} + PL(u^{(1)}, v^{(1)}, w^{(1)}) \otimes u^{(2)} \sqcup v^{(2)} \sqcup w^{(2)}. \quad (4)$$

Let $v \in T(V)$. Then:

$$\emptyset \bullet v = (\emptyset \sqcup \emptyset) \bullet v = (\emptyset \bullet v) \sqcup \emptyset + \emptyset \sqcup (\emptyset \bullet v) = 2\emptyset \bullet v,$$

so $\emptyset \bullet v = 0$ for any $v \in T(V)$. Hence, for any $v, w \in T(V)$, $PL(\emptyset, v, w) = 0$: by trilinearity of PL , we can assume that $\varepsilon(u) = 0$. In this case, (4) becomes:

$$\begin{aligned} \Delta(PL(u, v, w)) &= \emptyset \otimes PL(u, v, w) + PL(u, v^{(1)}, w^{(1)}) \otimes v^{(2)} \sqcup w^{(2)} \\ &\quad + PL(u', v^{(1)}, w^{(1)}) \otimes u'' \sqcup v^{(2)} \sqcup w^{(2)}. \end{aligned}$$

We assume that u, v, w are words of respective lengths k, l and n , with $k \geq 1$. Let us first prove that $PL(u, v, w) = 0$ if $l = 0$, or equivalently if $v = \emptyset$, by induction on n . If $n = 0$, then we can take $w = \emptyset$ and, obviously, $PL(u, \emptyset, \emptyset) = 0$. If $n \geq 1$, (4) becomes:

$$\begin{aligned} \Delta(PL(u, \emptyset, w)) &= \emptyset \otimes PL(u, v, w) + PL(u, \emptyset, w^{(1)}) \otimes w^{(2)} \\ &= \emptyset \otimes PL(u, v, w) + PL(u, \emptyset, w) \otimes \emptyset + PL(u, \emptyset, w') \otimes w''. \end{aligned}$$

By the induction hypothesis on n , $PL(u, \emptyset, w') = 0$, so $PL(u, \emptyset, w)$ is primitive, so belongs to V . As $\pi \circ PL = 0$, $PL(u, \emptyset, w) = 0$.

Hence, we can now assume that $l \geq 1$. By symmetry in v and w , we can also assume that $n \geq 1$. Let us now prove that $PL(u, v, w) = 0$ by induction on k . If $k = 0$, there is nothing more to prove. If $k \geq 1$, we proceed by induction on $l + n$. If $l + n \leq 1$, there is nothing more to prove. Otherwise, using both induction hypotheses, (4) becomes:

$$\Delta(PL(u, v, w)) = PL(u, v, w) \otimes \emptyset + \emptyset \otimes PL(u, v, w).$$

So $PL(u, v, w) \in V$. As $\pi \circ PL = 0$, $PL(u, v, w) = 0$. □

Consequently:

Proposition 8. Let $\varpi : T(V) \otimes T(V) \longrightarrow V$ be a linear map such that (2) and (3) are satisfied. The product \bullet defined by (1) makes $(T(V), \sqcup, \bullet, \Delta)$ a Com-PreLie bialgebra. We obtain in this way all the preLie products \bullet such that $(T(V), \sqcup, \bullet, \Delta)$ a Com-PreLie bialgebra. Moreover, for any $N \in \mathbb{Z}$, \bullet is homogeneous of degree N if, and only if:

$$\forall k, l \in \mathbb{N}, \quad k + l + N \neq 1 \implies \varpi(V^{\otimes k} \otimes V^{\otimes l}) = (0). \quad (5)$$

Remark 4. Let $\varpi : T(V) \otimes T(V) \longrightarrow V$, satisfying (5) for a given $N \in \mathbb{Z}$. Then:

1. (2) is satisfied if, and only if, for all $k, l, n \in \mathbb{N}$ such that $k + l + n = 1 - N$,

$$\forall u \in V^{\otimes k}, v \in V^{\otimes l}, w \in V^{\otimes n}, \quad \varpi((u \sqcup v) \otimes w) = \varepsilon(u)\varpi(v \otimes w) + \varepsilon(v)\varpi(u \otimes w).$$

2. (3) is satisfied if, and only if, for all $k, l, n \in \mathbb{N}$ such that $k + l + n = 1 - 2N$,

$$\begin{aligned} \forall u \in V^{\otimes k}, v \in V^{\otimes l}, w \in V^{\otimes n}, \quad & \varpi(u \bullet v \otimes w) - \varpi(u \otimes v \bullet w) \\ & = \varpi(u \bullet w \otimes v) - \varpi(u \otimes w \bullet v). \end{aligned}$$

Note that (2) is always satisfied if $u = \emptyset$ or $v = \emptyset$, that is to say if $k = 0$ or $l = 0$.

In the next paragraphs, we shall look at $N \geq 0$ and $N = -1$.

2.2 PreLie products of positive degree

Proposition 9. *Let f be a linear endomorphism of V . We define a product \bullet on $T(V)$ in the following way:*

$$\forall x_1, \dots, x_m, y_1, \dots, y_n \in V, \quad x_1 \dots x_m \bullet y_1 \dots y_n = \sum_{i=0}^n x_1 \dots x_{i-1} f(x_i)(x_{i+1} \dots x_m \sqcup y_1 \dots y_n). \quad (6)$$

Then $(T(V), \sqcup, \bullet, \Delta)$ is a Com-PreLie bialgebra denoted by $T(V, f)$. Conversely, if \bullet is a product on $T(V)$, homogeneous of degree $N \geq 0$, there exists a unique $f : V \rightarrow V$ such that $(T(V), \sqcup, \bullet, \Delta) = T(V, f)$.

Proof. We look for all possible ϖ , homogeneous of a certain degree $N \geq 0$, such that (2) and (3) are satisfied.

Let us consider such a ϖ . For any $k, l \in \mathbb{N}$, we denote by $\varpi_{k,l}$ the restriction of ϖ to $V^{\otimes k} \otimes V^{\otimes l}$. By (5), $\varpi_{k,l} = 0$ if $k + l \neq 1$. As (2) implies that $\varpi_{0,1} = 0$, the only possibly nonzero $\varpi_{k,l}$ is $\varpi_{1,0} : V \rightarrow V$, which we denote by f . Then (1) gives (6).

Let us consider any linear endomorphism f of V and consider ϖ such that the only nonzero component of ϖ is $\varpi_{1,0} = f$. Let us prove (2) for $u \in V^{\otimes k}, v \in V^{\otimes l}, w \in V^{\otimes n}$, with $k + l + n = 1 - N$. For all the possibilities for (k, l, n) , $0 \in \{k, l, n\}$, and the result is then obvious.

Let us prove (2) for $u \in V^{\otimes k}, v \in V^{\otimes l}, w \in V^{\otimes n}$, with $k + l + n = 1 - 2N$. We obtain two possibilities:

- $(k, l, n) = (0, 1, 0)$ or $(0, 0, 1)$. We can assume that $u = \emptyset$. As $\emptyset \bullet x = 0$ for any $x \in T(V)$, the result is obvious.
- $(k, l, n) = (1, 0, 0)$. We can assume that $v = w = \emptyset$, and the result is then obvious.

□

Remark 5. 1. If $N \geq 1$, necessarily $f = 0$, so $\bullet = 0$.

2. With the notation of Proposition 3, $f_{T(V,f)} = f$.

We obtain in this way the family of Com-PreLie bialgebras of [5], coming from a problem of composition of Fliess operators in Control Theory. Consequently, from [5]:

Corollary 10. *Let $k, l \geq 0$. We denote by $Sh(k, l)$ the set of (k, l) -shuffles, that is to say permutations $\sigma \in \mathfrak{S}_{k+l}$ such that:*

$$\sigma(1) < \dots < \sigma(k), \quad \sigma(k+1) < \dots < \sigma(k+l).$$

If $\sigma \in Sh(k, l)$, we put:

$$m_k(\sigma) = \max\{i \in [k] \mid \sigma(1) = 1, \dots, \sigma(i) = i\},$$

with the convention $m_k(\sigma) = 0$ if $\sigma(1) \neq 1$. Then, in $T(V, f)$, if $v_1, \dots, v_{k+l} \in V$:

$$v_1 \dots v_k \bullet v_{k+1} \dots v_{k+l} = \sum_{\sigma \in Sh(k,l)} \sum_{i=1}^{m_k(\sigma)} (Id^{\otimes(i-1)} \otimes f \otimes Id^{\otimes(k+l-i)})(v_{\sigma^{-1}(1)} \dots v_{\sigma^{-1}(k+l)}). \quad (7)$$

2.3 PreLie products of degree -1

Proposition 11. *Let $*$ and $\{-, -\}$ be two bilinear products on V such that:*

$$\begin{aligned} \forall x, y, z \in V, \quad (x * y) * z - x * (y * z) &= (x * z) * y - x * (z * y), \\ \{x, y\} &= -\{y, x\}, \\ x * \{y, z\} &= \{x * y, z\}, \\ \{x, y\} * z &= \{x * z, y\} + \{x, y * z\} + \{\{x, y\}, z\}. \end{aligned} \quad (8)$$

We define a product \bullet on $T(V)$ in the following way: for all $x_1, \dots, x_m, y_1, \dots, y_n \in V$,

$$\begin{aligned} x_1 \dots x_m \bullet y_1 \dots y_n &= \sum_{i=1}^n x_1 \dots x_{i-1} (x_i * y_1) (x_{i+1} \dots x_m \sqcup y_2 \dots y_n) \\ &\quad + \sum_{i=1}^{k-1} x_1 \dots x_{i-1} \{x_i, x_{i+1}\} (x_{i+2} \dots x_m \sqcup y_1 \dots y_n). \end{aligned} \quad (9)$$

Then $(T(V), \sqcup, \bullet, \Delta)$ is a Com-PreLie bialgebra, and we obtain in this way all the possible preLie products \bullet , homogeneous of degree -1 , such that $(T(V), \sqcup, \bullet, \Delta)$ is a Com-PreLie bialgebra.

Proof. Let us consider a linear map $\varpi : T(V) \otimes T(V) \rightarrow V$, satisfying (5) for $N = -1$. Denoting by $\varpi_{k,l} = \varpi|_{V^{\otimes k} \otimes V^{\otimes l}}$ for any k, l , the only possibly nonzero $\varpi_{k,l}$ are for $(k, l) = (2, 0)$, $(1, 1)$ and $(0, 2)$. For all $x, y \in V$, we put:

$$x * y = \varpi_{1,1}(x \otimes y), \quad \{x, y\} = \varpi_{2,0}(xy \otimes \emptyset).$$

(2) is equivalent to:

$$\begin{aligned} \forall w \in V^{\otimes 2}, \quad \varpi_{0,2}(\emptyset \otimes w) &= 0, \\ \forall x, y \in V, \quad \varpi_{2,0}((xy + yx) \otimes \emptyset) &= 0. \end{aligned}$$

Hence, we now assume that $\varpi_{0,2} = 0$, and we obtain that (2) is equivalent to (8)-2. The nullity of $\varpi_{0,2}$ and (1) give (9).

Let us now consider (3), with $u \in V^{\otimes k}$, $v \in V^{\otimes l}$, $w \in V^{\otimes n}$, $k + l + n = 1 - 2N = 3$. By symmetry between v and w , and by nullity of $\varpi_{0,l}$ for all l , we have to consider two cases:

- $k = l = n = 1$. We put $u = x$, $v = y$, $w = z$, with $x, y, z \in V$. Then (3) is equivalent to:

$$(x * y) * z - x * (y * z) = (x * z) * y - x * (z * y),$$

that is to say to (8)-1.

- $k = 1, l = 2, z = 0$. We put $u = x$, $v = yz$, $w = \emptyset$, with $x, y, z \in V$. Then (3) is equivalent to:

$$\{x * y, z\} - x * \{y, z\} = 0,$$

that is to say to (8)-3.

- $k = 2, l = 1, z = 0$. We put $u = xy$, $v = z$, $w = \emptyset$, with $x, y, z \in V$. Then (3) is equivalent to:

$$\{x * z, y\} + \{x, y * z\} + \{\{x, y\}, z\} = \{x, y\} * z,$$

that is to say to (8)-4.

We conclude with Proposition 8. □

Remark 6. 1. In particular, $*$ is a preLie product on V ; for all $x, y \in V$, $x \bullet y = x * y$.

2. If $x_1, \dots, x_m \in V$:

$$x_1 \dots x_m \bullet \emptyset = \sum_{i=1}^{m-1} x_1 \dots x_{i-1} \{x_i, x_{i+1}\} x_{i+2} \dots x_m.$$

Example 1. 1. If $*$ is a preLie product on V , we can take $\{-, -\} = 0$, and (8) is satisfied.

Using the classification of preLie algebras of dimension 2 over \mathcal{C} of [1], it is not difficult to show that if the dimension of V is 1 or 2, then necessarily $\{-, -\}$ is zero.

2. If $* = 0$, then (8) becomes:

$$\begin{aligned} \forall x, y \in V, & \quad \{x, y\} = -\{y, x\}, \\ \forall x, y, z \in V, & \quad \{\{x, y\}, z\} = 0, \end{aligned}$$

that is say $(V, \{-, -\})$ is a nilpotent Lie algebra, which nilpotency order is 2.

3. Here is a family of examples where both $*$ and $\{-, -\}$ are nonzero. Take V 3-dimensional, with basis (x, y, z) , a, b, c be scalars, and products given by the following arrays:

\bullet	x	y	z	$\{-, -\}$	x	y	z
x	x	y	z	x	0	$ay + bz$	$cy + (1 - a)z$
y	0	0	0	y	$-ay - bz$	0	0
z	0	0	0	z	$(a - 1)z - cy$	0	0

Then $(V, \bullet, \{-, -\})$ satisfies (8) if, and only if, $a^2 - a + bc = 0$, or equivalently:

$$(2a - 1)^2 + (b + c)^2 - (b - c)^2 = 1.$$

This equation defines a hyperboloid of one sheet.

3 Free Com-PreLie algebras and quotients

3.1 Description of free Com-PreLie algebras

We described in [5] free Com-PreLie algebras in terms of decorated rooted partitioned trees. We now work with free unitary Com-PreLie algebras.

Definition 12. 1. A partitioned forest is a pair (F, I) such that:

- (a) F is a rooted forest (the edges of F being oriented from the roots to the leaves). The set of its vertices is denoted by $V(F)$.
- (b) I is a partition of the vertices of F with the following condition: if x, y are two vertices of F which are in the same part of I , then either they are both roots, or they have the same direct ascendant.

The parts of the partition are called blocks.

- 2. We shall say that a partitioned forest F is a partitioned tree if all the roots are in the same block. Note that in this case, one of the blocks of F is the set of roots of F . By convention, the empty forest \emptyset is considered as a partitioned tree.
- 3. Let \mathcal{D} be a set. A partitioned tree decorated by \mathcal{D} is a triple (T, I, d) , where (T, I) is a partitioned tree and d is a map from the set of vertices of T into \mathcal{D} . For any vertex x of T , $d(x)$ is called the decoration of x .

Proof. Obviously, (A, \cdot) is a commutative, unitary associative algebra. Let us prove the PreLie identity for $x, y, z \in A_+ \sqcup \{\emptyset\}$.

- If $x = \emptyset$, then $x \bullet (y \bullet z) = (x \bullet y) \bullet z = x \bullet (z \bullet y) = (x \bullet z) \bullet y = 0$. We now assume that $x \in A_+$.
- If $y = z = \emptyset$, then obviously the PreLie identity is satisfied.
- If $y = \emptyset$ and $z \in A_+$, then:

$$\begin{aligned} x \bullet (y \bullet z) &= 0, & (x \bullet y) \bullet z &= f(x) \bullet y, \\ x \bullet (z \bullet y) &= x \bullet f(z), & (x \bullet z) \bullet y &= f(x \bullet z). \end{aligned}$$

As f is a derivation for \bullet , the PreLie identity is satisfied. By symmetry, it is also true if $y \in A_+$ and $z = \emptyset$.

Let us now prove the Leibniz identity for $x, y, z \in A_+ \sqcup \{\emptyset\}$. It is obviously satisfied if $x = \emptyset$ or $y = \emptyset$; we assume that $x, y \in A_+$. If $z = \emptyset$, then:

$$(x \cdot y) \bullet z = f(x \cdot y), \quad (x \bullet z) \cdot y = f(x) \cdot y, \quad x \cdot (y \bullet z) = x \cdot f(y).$$

As f is a derivation for \cdot , the Leibniz identity is satisfied. \square

Proposition 15. *Let $UCP(\mathcal{D})$ be the vector space generated by $UPT(\mathcal{D})$. We extend \cdot by bilinearity and the PreLie product \bullet is defined by:*

$$\forall T, T' \in UPT(\mathcal{D}), \quad T \bullet T' = \begin{cases} \sum_{s \in V(t)} T \bullet_{s,*} T' & \text{if } t \neq \emptyset, \\ \sum_{s \in V(t)} T[+1]_s & \text{if } t = \emptyset. \end{cases}$$

Then $UCP(\mathcal{D})$ is the free unitary Com-PreLie algebra generated by the elements $\bullet_{(0,d)}$, $d \in \mathcal{D}$.

Proof. We denote by $UCP_+(\mathcal{D})$ the subspace of $UCP(\mathcal{D})$ generated by nonempty trees. By proposition 18 in [5], this is the free Com-PreLie algebra generated by the elements $\bullet_{(k,d)}$, $k \in \mathbb{N}$, $d \in \mathcal{D}$. We define a map $f : UCP_+(\mathcal{D}) \rightarrow UCP_+(\mathcal{D})$ by:

$$\forall T \in UPT(\mathcal{D}) \setminus \{\emptyset\}, \quad f(T) = \sum_{s \in V(t)} T[+1]_s.$$

This is a derivation for both \cdot and \bullet ; by lemma 14, $UCP(\mathcal{D})$ is a unitary Com-PreLie algebra.

Observe that for all $d \in \mathcal{D}$, $k \in \mathbb{N}$:

$$\bullet_{(0,d)} \bullet^{\times k} = \bullet_{(k,d)}.$$

Let A be a unitary Com-PreLie algebra and, for all $d \in \mathcal{D}$, let $a_d \in A$. By proposition 18 in [5], we define a unique Com-PreLie algebra morphism:

$$\theta : \begin{cases} UCP_+(\mathcal{D}) & \longrightarrow A \\ \bullet_{(k,d)} & \longrightarrow a_d \times 1_A^{\times k}. \end{cases}$$

We extend it to $UCP(\mathcal{D})$ by sending \emptyset to 1_A , and we obtain in this way a unitary Com-PreLie algebra from $UCP(\mathcal{D})$ to A , sending $\bullet_{(0,d)}$ to a_d for any $d \in \mathcal{D}$. This morphism is clearly unique. \square

Example 4. Let $i, j, k \in \mathbb{N}$ and $d, e, f \in \mathcal{D}$.

$$\begin{aligned}
\bullet_{(i,d)} \bullet_{(j,e)} &= \mathfrak{!}_{(i,d)}^{(j,e)}, \\
\bullet_{(i,d)} \bullet_{(j,e)} \dashrightarrow (k,f) &= \overset{(j,e)}{\mathfrak{V}}_{(i,d)}^{(k,f)}, \\
\bullet_{(i,d)} \bullet \mathfrak{!}_{(j,e)}^{(k,f)} &= \mathfrak{!}_{(i,d)}^{(k,f)}, \\
\mathfrak{!}_{(i,d)}^{(j,e)} \bullet_{(k,f)} &= \mathfrak{!}_{(i,d)}^{(k,f)} + \overset{(j,e)}{\mathfrak{V}}_{(i,d)}^{(k,f)}, \\
\bullet_{(i,d)} \bullet \emptyset &= \bullet_{(i+1,d)}, \\
\mathfrak{!}_{(i,d)}^{(j,e)} \bullet \emptyset &= \mathfrak{!}_{(i+1,d)}^{(j,e)} + \mathfrak{!}_{(i,d)}^{(j+1,e)}, \\
\overset{(j,e)}{\mathfrak{V}}_{(i,d)}^{(k,f)} \bullet \emptyset &= \overset{(j,e)}{\mathfrak{V}}_{(i+1,d)}^{(k,f)} + \overset{(j+1,e)}{\mathfrak{V}}_{(i,d)}^{(k,f)} + \overset{(j,e)}{\mathfrak{V}}_{(i,d)}^{(k+1,f)}.
\end{aligned}$$

3.2 Quotients of $UCP(\mathcal{D})$

Proposition 16. We put $V_0 = \text{Vect}(\bullet_{(0,d)}, d \in \mathcal{D})$, identified with $\text{Vect}(\bullet_d, d \in \mathcal{D})$. Let $f : V_0 \rightarrow V_0$ be any linear map. We consider the Com-PreLie ideal I_f of $UCP(\mathcal{D})$ generated by the elements $\bullet_{(1,d)} - f(\bullet_{(0,d)})$, $d \in \mathcal{D}$.

1. We denote by $\mathcal{UPT}'(\mathcal{D})$ the set of trees $T \in \mathcal{UPT}(\mathcal{D})$ such that for any vertex s of T , the decoration of s is of the form $(0, d)$, with $d \in \mathcal{D}$. It is trivially identified with $\mathcal{PT}(\mathcal{D})$. Then the family $(T + I_f)_{T \in \mathcal{UPT}'(\mathcal{D})}$ is a basis of $UCP(\mathcal{D})/I_f$.
2. In $UCP(\mathcal{D})/I_f$, for any $d \in \mathcal{D}$, $(\bullet_d + I_f) \bullet \emptyset = f(\bullet_d)$.

Proof. *First step.* We fix $d \in \mathcal{D}$. Let us first prove that for all $k \geq 0$:

$$\bullet_{(k,d)} + I_f = f^k(\bullet_{(0,d)}) + I_f.$$

It is obvious if $k = 0, 1$. Let us assume the result at rank $k - 1$. We put $f(\bullet_{(0,d)}) = \sum a_e \bullet_{(0,e)}$. Then:

$$\begin{aligned}
\bullet_{(k,d)} + I_f &= \bullet_{(1,d)} \bullet \emptyset^{\times(k-1)} + I_f \\
&= \sum a_e \bullet_{(0,e)} \bullet \emptyset^{\times(k-1)} + I_f \\
&= \sum a_e f^{k-1}(\bullet_{(0,e)}) + I_f \\
&= f^k(\bullet_{(0,d)}) + I_f,
\end{aligned}$$

so the result holds for all k .

Second step. Let $T \in \mathcal{UPT}(\mathcal{D})$; let us prove that there exists $x \in \text{Vect}(\mathcal{UPT}'(\mathcal{D}))$, such that $T + I_f = x + I_f$. We proceed by induction on $|T|$. If $|T| = 0$, then $t = \emptyset$ and we can take $x = T$. If $|T| = 1$, then $T = \bullet_{(k,d)}$ and we can take, by the first step, $x = f^k(\bullet_{(0,d)})$. Let us assume the result at all ranks $< |T|$. If T has several roots, we can write $T = T_1 \cdot T_2$, with $|T_1|, |T_2| < |T|$. Hence, there exists $x_i \in \text{Vect}(\mathcal{UPT}'(\mathcal{D}))$, such that $T_i + I_f = x_i + I_f$ for all $i \in [2]$, and we take $x = x_1 \cdot x_2$. Otherwise, we can write:

$$T = \bullet_{(k,d)} \bullet T_1 \times \dots \times T_k,$$

where $T_1, \dots, T_k \in \mathcal{UPT}(\mathcal{D})$. By the induction hypothesis, there exists $x_i \in \text{Vect}(\mathcal{UPT}'(\mathcal{D}))$ such that $T_i + I_f = x_i + I_f$ for all $i \in [k]$. We then take $x = f^k(\bullet_{(0,d)}) \bullet x_1 \times \dots \times x_k$.

Third step. We give $CP_+(\mathcal{D}) = \text{Vect}(\mathcal{PT}(\mathcal{D}) \setminus \{\emptyset\})$ a Com-PreLie structure by:

$$\forall T, T' \in \mathcal{PT}(\mathcal{D}) \setminus \{\emptyset\}, T \bullet T' = \sum_{s \in V(t)} T \bullet_{s,*} T'.$$

We consider the map:

$$F : \begin{cases} CP_+(\mathcal{D}) & \longrightarrow CP_+(\mathcal{D}) \\ T & \longrightarrow \sum_{s \in V(T)} f_s(T), \end{cases}$$

where, $f_s(T)$ is the linear span of decorated partitioned trees obtained by replacing the decoration d_s of s by $f(d_s)$, the trees being considered as linear in any of their decorations. This is a derivation for both \cdot and \bullet , so by lemma 14, $CP(\mathcal{D})$ inherits a unitary Com-PreLie structure such that for any $d \in \mathcal{D}$:

$$\cdot_d \bullet \emptyset = f(\cdot_d).$$

By the universal property of $UCP(\mathcal{D})$, there exists a unique unitary Com-PreLie algebra structure $\phi : UCP(\mathcal{D}) \longrightarrow CP(\mathcal{D})$, such that $\phi(\cdot_{(0,d)}) = \cdot_d$ for any $d \in \mathcal{D}$. Then $\phi(\cdot_{(1,d)}) = f(\cdot_d) = \phi(f(\cdot_{(0,d)}))$ for any $d \in \mathcal{D}$, so ϕ induces a morphism $\bar{\phi} : UCP(\mathcal{D})/I_f \longrightarrow CP(\mathcal{D})$. It is not difficult to prove that for any $T \in \mathcal{UP}\mathcal{T}'(\mathcal{D})$, $\phi(T) = T$. As the family $\mathcal{PT}(\mathcal{D})$ is a basis of $CP(\mathcal{D})$, the family $(T + I_f)_{T \in \mathcal{UP}\mathcal{T}'(\mathcal{D})}$ is linearly independent in $UCP(\mathcal{D})/I_f$. By the second step, it is a basis. \square

Example 5. We choose $f = Id_{V_0}$. The product in $UCP(\mathcal{D})/I_{Id_{V_0}}$ of two elements is given by the combinatorial product \cdot . If $T, T' \in \mathcal{PT}(\mathcal{D})$ and $T' \neq \emptyset$, $T \bullet T'$ is the sum of all graftings of T' over T . Moreover:

$$T \bullet \emptyset = |T|T.$$

Hence, we now consider $CP(\mathcal{D})$, augmented by an unit \emptyset , as a unitary Com-PreLie algebra.

Proposition 17. *Let J be the Com-PreLie ideal of $CP(\mathcal{D})$ generated by the elements $\cdot_d \bullet (F_1 \times F_2) - \cdot_d \bullet (F_1 \cdot F_2)$, with $d \in \mathcal{D}$ and $F_1, F_2 \in \mathcal{PT}(\mathcal{D})$.*

1. *Let T and T' be two elements of $\mathcal{PT}(\mathcal{D})$ which are equal as decorated rooted forests. Then $T + J = T' + J$. Consequently, if F is a decorated rooted forest, the element $T' + J$ does not depend of the choice of $T' \in \mathcal{UP}\mathcal{T}(\mathcal{D})$ such that $T' = F$ as a decorated rooted forest. This element is identified with F .*
2. *The set of decorated rooted forests is a basis of $UCP(\mathcal{D})/J$.*

$CP(\mathcal{D})/J$ is then, as an algebra, identified with the Connes-Kreimer algebra $H_{CK}^{\mathcal{D}}$ of decorated rooted trees [3, 4], which is in this way a unitary Com-PreLie algebra.

Proof. 1. *First step.* Let us show that for any $x_1, \dots, x_n \in UCP(\mathcal{D})$, $\cdot_d \bullet (x_1 \times \dots \times x_n) + J = \cdot_d \bullet (x_1 \cdot \dots \cdot x_n) + J$ by induction on n . It is obvious if $n = 1$, and it comes from the definition of J if $n = 2$. Let us assume the result at rank $n - 1$.

$$\begin{aligned} & \cdot_d \bullet (x_1 \times \dots \times x_n) + J \\ &= (\cdot_d \bullet (x_1 \times \dots \times x_{n-1})) \bullet x_n - \sum_{i=1}^{n-1} \cdot_d \bullet (x_1 \times \dots \times (x_i \bullet x_n) \times \dots \times x_{n-1}) + J \\ &= (\cdot_d \bullet (x_1 \cdot \dots \cdot x_{n-1})) \bullet x_n - \sum_{i=1}^{n-1} \cdot_d \bullet (x_1 \cdot \dots \cdot (x_i \bullet x_n) \cdot \dots \cdot x_{n-1}) + J \\ &= (\cdot_d \bullet (x_1 \cdot \dots \cdot x_{n-1})) \bullet x_n - \cdot_d \bullet ((x_1 \cdot \dots \cdot x_{n-1}) \bullet x_n) + J \\ &= \cdot_d \bullet ((x_1 \cdot \dots \cdot x_{n-1}) \times x_n) + J \\ &= \cdot_d \bullet (x_1 \cdot \dots \cdot x_{n-1} \cdot x_n) + J. \end{aligned}$$

So the result holds for all n .

Second step. Let $F, G \in \mathcal{PT}(\mathcal{D})$, such that the underlying rooted decorated forests are equal. Let us prove that $F + J = G + J$ by induction on $n = |F| = |G|$. If $n = 0$, $F = G = 1$ and it is obvious. If $n = 1$, $F = G = \cdot_d$ and it is obvious. Let us assume the result at all ranks $< n$.

First case. If F has $k \geq 2$ roots, we can write $F = T_1 \cdot \dots \cdot T_k$ and $G = T'_1 \cdot \dots \cdot T'_k$, such that, for all $i \in [k]$, T_i and T'_i have the same underlying decorated rooted forest; By the induction hypothesis, $T_i + J = T'_i + J$ for all i , so $F + J = G + J$.

Second case. Let us assume that F has only one root. We can write $F = \cdot_d \bullet (F_1 \times \dots \times F_k)$ and $G = \cdot_d \bullet (G_1 \times \dots \times G_l)$. Then $F_1 \cdot \dots \cdot F_k$ and $G_1 \cdot \dots \cdot G_l$ have the same underlying decorated forest; by the induction hypothesis, $F_1 \cdot \dots \cdot F_k + J = G_1 \cdot \dots \cdot G_l + J$, so $\cdot_d \bullet (F_1 \cdot \dots \cdot F_k) + J = \cdot_d \bullet (G_1 \cdot \dots \cdot G_l) + J$. By the first step:

$$F + J = \cdot_d \bullet (F_1 \cdot \dots \cdot F_k) + J = \cdot_d \bullet (G_1 \cdot \dots \cdot G_l) + J = G + J.$$

2. The set $\mathcal{RF}(\mathcal{D})$ of rooted forests linearly spans $CP(\mathcal{D})/J$ by the first point. Let J' be the subspace of $CP(\mathcal{D})$ generated by the differences of elements of $\mathcal{PT}(\mathcal{D})$ with the same underlying decorated forest. It is clearly a Com-PreLie ideal, and $\mathcal{RF}(\mathcal{D})$ is a basis of $CP(\mathcal{D})/J'$. Moreover, for all $F_1, F_2 \in \mathcal{PT}(\mathcal{D})$, $\cdot_d \bullet (F_1 \times F_2) + J' = \cdot_s \bullet (F_1 \cdot F_2) + J'$, as the underlying forests of $\cdot_d \bullet (F_1 \times F_2)$ and $\cdot_s \bullet (F_1 \cdot F_2)$ are equal. Consequently, there exists a Com-PreLie morphism from $CP(\mathcal{D})/J$ to $CP(\mathcal{D})/J'$, sending any element of $\mathcal{RF}(\mathcal{D})$ over itself. As the elements of $RF(\mathcal{D})$ are linearly independent in $CP(\mathcal{D})/J'$, they also are in $CP(\mathcal{D})/J$. \square

3.3 PreLie structure of $UCP(\mathcal{D})$ and $CP(\mathcal{D})$

Let us now consider $UCP(\mathcal{D})$ and $CP(\mathcal{D})$ as PreLie algebras. Their augmentation ideals are respectively denoted by $UCP_+(\mathcal{D})$ and $CP_+(\mathcal{D})$. Note that, as a PreLie algebra, $UCP_+(\mathcal{D}) = CP_+(\mathbb{N} \times \mathcal{D})$.

Let \mathcal{D} be any set, and let $T \in \mathcal{PT}(\mathcal{D})$. Then T can be written as:

$$T = (\cdot_{d_1} \bullet (T_{1,1} \times \dots \times T_{i,s_1})) \cdot \dots \cdot (\cdot_{d_k} \bullet (T_{k,1} \times \dots \times T_{k,s_k})),$$

where $d_1, \dots, d_k \in \mathcal{D}$ and the $T_{i,j}$'s are nonempty elements of $\mathcal{PT}(\mathcal{D})$. We shortly denote this as:

$$T = B_{d_1, \dots, d_k}(T_{1,1} \dots T_{1,s_1}; \dots; T_{k,1} \dots T_{k,s_k}).$$

The set of partitioned subtrees $T_{i,j}$ of T is denoted by $st(T)$.

Proposition 18. *Let \mathcal{D} be any set. One defines a coproduct δ on $CP_+(\mathcal{D})$ by:*

$$\forall T \in \mathcal{PT}(\mathcal{D}), \quad \delta(T) = \sum_{T' \in st(T)} T \setminus T' \otimes T.$$

Then, as a PreLie algebra, $CP_+(\mathcal{D})$ is freely generated by $Ker(\delta)$.

Proof. In other words, for any $T \in \mathcal{PT}(\mathcal{D})$, writing

$$T = B_{d_1, \dots, d_k}(T_{1,1} \dots T_{1,s_1}; \dots; T_{k,1} \dots T_{k,s_k}).$$

we have:

$$\delta(T) = \sum_{i=1}^s \sum_{j=1}^{s_i} B_{d_1, \dots, d_k}(T_{1,1} \dots T_{1,s_1}; \dots; T_{i,1} \dots \widehat{T_{i,j}} \dots T_{i,s_i}; \dots; T_{k,1} \dots T_{k,s_k}) \otimes T_{i,j}.$$

This immediately implies that δ is permutative [9]:

$$(\delta \otimes Id) \circ \delta = (23) \cdot (\delta \otimes Id) \circ \delta.$$

Moreover, for any $x, y \in \mathcal{PT}_+(\mathcal{D})$, using Sweedler's notation $\delta(x) = x^{(1)} \otimes x^{(2)}$, we obtain:

$$\delta(x \cdot y) = x^{(1)} \cdot y \otimes x^{(2)} + x \cdot y^{(1)} \otimes y^{(2)}.$$

For any partitioned tree $T \in \mathcal{PT}(\mathcal{D})$, we denote by $r(T)$ the number of roots of T and we put $d(T) = r(T)T$. The map d is linearly extended as an endomorphism of $\mathcal{PT}(\mathcal{D})$. As the product \cdot is homogeneous for the number of roots, d is a derivation of the algebra $(CP(\mathcal{D}), \cdot)$. Let us prove that for any $x, y \in CP_+(\mathcal{D})$:

$$\delta(x \bullet y) = d(x) \otimes y + x^{(1)} \bullet y \otimes x^{(2)} + x^{(1)} \otimes x^{(2)} \bullet y.$$

We denote by A the set of elements $x \in CP_+(\mathcal{D})$, such that for any $y \in CP_+(\mathcal{D})$, the preceding equality holds. If $x_1, x_2 \in A$, then for any $y \in CP_+(\mathcal{D})$:

$$\begin{aligned} \delta((x_1 \cdot x_2) \bullet y) &= \delta((x_1 \bullet y) \cdot x_2) + \delta(x_1 \cdot (x_2 \bullet y)) \\ &= (x_1 \bullet y)^{(1)} \cdot x_2 \otimes (x_1 \bullet y)^{(2)} + (x_1 \bullet y) \cdot x_2^{(1)} \otimes x_2^{(2)} \\ &\quad + x_1^{(1)} \cdot (x_2 \bullet y) \otimes x_1^{(2)} + x_1 \cdot (x_2 \bullet y)^{(1)} \otimes (x_2 \bullet y)^{(2)} \\ &= d(x_1) \cdot x_2 \otimes y + (x_1^{(1)} \bullet y) \cdot x_2 \otimes x_1^{(1)} + x_1^{(1)} \cdot x_2 \otimes x_1^{(2)} \bullet y \\ &\quad + (x_1 \bullet y) \cdot x_2^{(1)} \otimes x_2^{(2)} + x_1^{(1)} \cdot (x_2 \bullet y) \otimes x_1^{(2)} \\ &\quad + x_1 \cdot d(x_2) \otimes y + x_1 \cdot (x_2^{(1)} \bullet y) \otimes x_2^{(2)} + x_1 \cdot x_2^{(1)} \otimes x_2^{(2)} \bullet y \\ &= d(x_1 \cdot x_2) \otimes y + (x_1^{(1)} \cdot x_2) \bullet y \otimes x_1^{(2)} + (x_1 \cdot x_2^{(1)}) \bullet y \otimes x_2^{(2)} \\ &\quad + (x_1 \cdot x_2)^{(1)} \otimes (x_1 \cdot x_2)^{(2)} \bullet y \\ &= d(x_1 \cdot x_2) \otimes y + (x_1 \cdot x_2)^{(1)} \bullet y \otimes (x_1 \cdot x_2)^{(2)} + (x_1 \cdot x_2)^{(1)} \otimes (x_1 \cdot x_2)^{(2)} \bullet y. \end{aligned}$$

So $x_1 \cdot x_2 \in A$.

Let $d \in \mathcal{D}$. Note that $\delta(\cdot_a) = 0$. Moreover, for any $y \in CP_+(\mathcal{D})$:

$$\delta(\cdot_a \bullet y) = \delta(B_d(y)) = \cdot_a \otimes y,$$

so $\cdot_a \in A$. Let $T_1, \dots, T_k \in \mathcal{PT}(\mathcal{D})$, nonempty. We consider $x = B_d(T_1 \dots T_k)$. For any $y \in CP_+(\mathcal{D})$:

$$\begin{aligned} \delta(x \bullet y) &= \delta(B_d(T_1 \dots T_k y)) + \sum_{j=1}^k \delta(B_d(T_1 \dots (T_j \bullet y) \dots T_k)) \\ &= B_d(T_1 \dots T_k) \otimes y + \sum_{i=1}^k D_d(T_1 \dots \widehat{T}_i \dots T_k y) \otimes T_i \\ &\quad + \sum_{i=1}^k \sum_{j \neq i} B_d(T_1 \dots \widehat{T}_i \dots (T_j \bullet y) \dots T_k) \otimes T_i + \sum_{i=1}^k B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \bullet y \\ &= d(x) \otimes y + \sum_{i=1}^k B_d(T_1 \dots \widehat{T}_i \dots T_k) \bullet y \otimes T_i + \sum_{i=1}^k B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \bullet y \\ &= d(x) \otimes y + x^{(1)} \bullet y \otimes x^{(2)} + x^{(1)} \otimes x^{(2)} \bullet y. \end{aligned}$$

Hence, $x \in A$. As A is stable under \cdot and contains any partitioned tree with one root, $A = CP_+(\mathcal{D})$.

For any nonempty partitioned tree $T \in \mathcal{PT}(\mathcal{D})$, we put $\delta'(T) = \frac{1}{r(T)}\delta(T)$. Then:

$$(\delta' \otimes Id) \circ \delta'(T) = \frac{1}{r(T)^2}(\delta \otimes Id) \circ \delta(T),$$

so δ' is also permutative; moreover, for any $x, y \in CP_+(\mathcal{D})$:

$$\delta'(x \bullet y) = x \otimes y + x^{(1)} \bullet y \otimes x^{(2)} + x^{(1)} \otimes x^{(2)} \bullet y.$$

By Livernet's rigidity theorem [9], the PreLie algebra $CP_+(\mathcal{D})$ is freely generated by $Ker(\delta')$. For any integer n , we denote by $CP_n(\mathcal{D})$ the subspace of $CP(\mathcal{D})$ generated by trees T such that $r(T) = n$. Then, for all n , $\delta(CP_n(\mathcal{D})) \subseteq CP_n(\mathcal{D}) \otimes CP_+(\mathcal{D})$, and $\delta|_{CP_n(\mathcal{D})} = n\delta'|_{CP_n(\mathcal{D})}$. This implies that $Ker(\delta) = Ker(\delta')$. \square

Lemma 19. *In $CP_+(\mathcal{D})$ or $UCP_+(\mathcal{D})$, $Ker(\delta) \bullet \emptyset \subseteq Ker(\delta)$.*

Proof. Let us work in $UCP_+(\mathcal{D})$. Let us prove that for any $x \in UCP_+(\mathcal{D})$:

$$\delta(x \bullet \emptyset) = x^{(1)} \bullet \emptyset \otimes x^{(2)} + x^{(1)} \otimes x^{(2)} \bullet \emptyset.$$

We denote by A the subspace of elements $x \in UCP_+(\mathcal{D})$ such that this holds. If $x_1, x_2 \in A$, then:

$$\begin{aligned} \delta((x_1 \cdot x_2) \bullet \emptyset) &= \delta((x_1 \bullet \emptyset) \cdot x_2) + \delta(x_1 \cdot (x_2 \bullet \emptyset)) \\ &= (x_1^{(1)} \bullet \emptyset) \cdot x_2 \otimes x^{(1)} + x_1^{(1)} \cdot x_2 \otimes x_1^{(2)} \bullet \emptyset + (x_1 \bullet \emptyset) \cdot x_2^{(1)} \otimes x_2^{(2)} \\ &\quad + x_1 \cdot (x_2^{(1)} \bullet \emptyset) \otimes x_2^{(2)} + x_1 \cdot x_2^{(1)} \otimes x_2^{(2)} \bullet \emptyset + x_1^{(1)} \cdot (x_2 \bullet \emptyset) \otimes x_1^{(2)} \\ &= (x_1^{(1)} \cdot x_2) \bullet \emptyset \otimes x_1^{(2)} + x_1^{(1)} \cdot x_2 \otimes x_1^{(2)} \bullet \emptyset \\ &\quad + (x_1 \cdot x_2^{(1)}) \bullet \emptyset \otimes x_2^{(2)} + x_1 \cdot x_2^{(1)} \otimes x_2^{(2)} \bullet \emptyset \\ &= (x_1 \cdot x_2)^{(1)} \bullet \emptyset \otimes (x_1 \cdot x_2)^{(2)} + (x_1 \cdot x_2)^{(1)} \otimes (x_1 \cdot x_2)^{(2)} \bullet \emptyset, \end{aligned}$$

so $x_1 \cdot x_2 \in A$. If $d \in D$ and $T_1, \dots, T_k \in \mathcal{PT}(\mathcal{D})$, nonempty, if $x = B_d(T_1 \dots T_k)$:

$$\begin{aligned} \delta(x \bullet \emptyset) &= \delta(B_{d+1}(T_1 \dots T_k)) + \sum_{i=1}^k \delta(B_d(T_1 \dots (T_i \bullet \emptyset) \dots T_k)) \\ &= \sum_{i=1}^k B_{d+1}(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i + \sum_{j=1}^k \sum_{i \neq j} B_d(T_1 \dots (T_j \bullet \emptyset) \dots \widehat{T}_i \dots T_k) \otimes T_i \\ &\quad + \sum_{i=1}^k B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \bullet \emptyset \\ &= \sum_{i=1}^k B_d(T_1 \dots \widehat{T}_i \dots T_k) \bullet \emptyset \otimes T_i + \sum_{i=1}^k B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \bullet \emptyset \\ &= x^{(1)} \bullet \emptyset \otimes x^{(2)} + x^{(1)} \otimes x^{(2)} \bullet \emptyset, \end{aligned}$$

so $x \in A$. Hence, $A = UCP_+(\mathcal{D})$. Consequently, if $x \in Ker(\delta)$, then $x \bullet \emptyset \in Ker(\delta)$. The proof is immediate for $CP_+(\mathcal{D})$, as for any tree $T \in \mathcal{PT}(\mathcal{D})$, $T \bullet \emptyset = |T|T$. \square

We denote by ϕ the endomorphism of $Ker(\delta)$ defined by $\phi(x) = x \bullet \emptyset$.

Corollary 20. *The PreLie algebra $UCP(\mathcal{D})$, respectively $CP(\mathcal{D})$, is generated by $Ker(\delta) \oplus (\emptyset)$, with the relations:*

$$\begin{aligned} \forall x \in Ker(\delta), \quad & \emptyset \bullet \emptyset = 0, \\ & \emptyset \bullet x = 0, \quad x \bullet \emptyset = \phi(x). \end{aligned}$$

Remark 7. We give $CP(\mathcal{D})$ a graduation by putting the elements of \mathcal{D} homogeneous of degree 1. A manipulation of formal series allows to compute the dimensions of the homogeneous components of $Ker(\delta)$, if $|D| = d$:

$$\begin{aligned} \dim(Ker(\delta)_1) &= d, \\ \dim(Ker(\delta)_2) &= \frac{d(d+1)}{2}, \\ \dim(Ker(\delta)_3) &= \frac{d(2d^2+1)}{3}, \\ \dim(Ker(\delta)_4) &= \frac{d(11d^3+2d^2+d+2)}{8}, \\ \dim(Ker(\delta)_5) &= \frac{d(203d^4+60d^3-5d^2-30d+12)}{60}, \\ \dim(Ker(\delta)_6) &= \frac{d(220d^5+89d^4+16d^3+3d^2+4d+4)}{24}. \end{aligned}$$

4 Bialgebra structures on free Com-PreLie algebras

4.1 Tensor product of Com-PreLie algebras

Lemma 21. *Let A_1, A_2 be two Com-PreLie algebras and let $\varepsilon : A_1 \rightarrow \mathbb{K}$ such that:*

$$\forall a, b \in A_1, \varepsilon(a \bullet b) = \varepsilon(b \bullet a).$$

Then $A_1 \otimes A_2$ is a Com-PreLie algebra, with the products defined by:

$$\begin{aligned} (a_1 \otimes a_2)(b_1 \otimes b_2) &= a_1 b_1 \otimes a_2 b_2, \\ (a_1 \otimes a_2) \bullet_\varepsilon (b_1 \otimes b_2) &= a_1 \bullet b_1 \otimes a_2 b_2 + \varepsilon(b_1) a_1 \otimes a_2 \bullet b_2. \end{aligned}$$

Proof. $A_1 \otimes A_2$ is obviously an associative and commutative algebra, with unit $1 \otimes 1$. We take $A = a_1 \otimes a_2, B = b_1 \otimes b_2, C = c_1 \otimes c_2 \in A_1 \otimes A_2$. Let us prove the PreLie identity.

$$\begin{aligned} (A \bullet_\varepsilon B) \bullet_\varepsilon C - A \bullet_\varepsilon (B \bullet_\varepsilon C) &= (a_1 \bullet b_1) \bullet c_1 \otimes a_2 b_2 c_2 + \varepsilon(c_1) a_1 \bullet b_1 \otimes (a_2 b_2) \bullet c_2 \\ &\quad + \varepsilon(b_1) a_1 \bullet c_1 \otimes (a_2 \bullet b_2) c_2 + \varepsilon(b_1) \varepsilon(c_1) a_1 \otimes (a_2 b \bullet_2) \bullet c_2 \\ &\quad - a_1 \bullet (b_1 \bullet c_1) \otimes a_2 b_2 c_2 - \varepsilon(c_1) a_1 \bullet b_1 \otimes a_2 (b_2 \bullet c_2) \\ &\quad - \varepsilon(c_1) \varepsilon(b_1) a_1 \otimes a_2 \bullet (b_2 \bullet c_2) - \varepsilon(b_1 \bullet c_1) a_1 \otimes a_2 \bullet (b_2 c_2) \\ &= ((a_1 \bullet b_1) \bullet c_1 - a_1 \bullet (b_1 \bullet c_1)) \otimes a_2 b_2 c_2 \\ &\quad + \varepsilon(b_1) \varepsilon(c_1) a_1 \otimes ((a_2 \bullet b_2) \bullet c_2 - a_2 \bullet (b_2 \bullet c_2)) \\ &\quad + \varepsilon(c_1) a_1 \bullet b_1 \otimes (a_2 \bullet c_2) b_2 + \varepsilon(b_1) a_1 \bullet c_1 \otimes (a_2 \bullet b_2) c_2 \\ &\quad - \varepsilon(b_1 \bullet c_1) a_1 \otimes a_2 \bullet (b_2 c_2). \end{aligned}$$

As A_1 and A_2 are PreLie, the first and second lines of the last equality are symmetric in B and C ; the third line is obviously symmetric in B and C ; as m is commutative and by the hypothesis on ε , the last line also is. So \bullet_ε is PreLie.

$$\begin{aligned} (AB) \bullet C &= (a_1 b_1) \bullet c_1 \otimes a_2 b_2 c_2 + \varepsilon(c_1) a_1 b_1 \otimes (a_2 b_2) \bullet c_2 \\ &= ((a_1 \bullet c_1) b_1 + a_1 (b_1 \bullet c_1)) \otimes a_2 b_2 c_2 + \varepsilon(c_1) a_1 b_1 \otimes ((a_2 \bullet c_2) b_2 + a_2 (b_2 \bullet c_2)) \\ &= (a_1 \bullet c_1 \otimes a_2 c_2 + \varepsilon(c_1) a_1 \otimes a_2 \bullet c_2) (b_1 \otimes b_2) \\ &\quad + (a_1 \otimes a_2) (b_1 \bullet c_1 \otimes b_2 c_2 + \varepsilon(c_1) b_1 \otimes b_2 \bullet c_2) \\ &= (A \bullet C) B + A (B \bullet C). \end{aligned}$$

So $A_1 \otimes A_2$ is Com-PreLie. □

Remark 8. Consequently, if (A, m, \bullet, Δ) is a Com-PreLie bialgebra, with counit ε , then Δ is a morphism of Com-PreLie algebras from (A, m, \bullet) to $(A \otimes A, m, \bullet_\varepsilon)$. Indeed, for all $a, b \in A$, $\varepsilon(a \bullet b) = \varepsilon(b \bullet a) = 0$ and:

$$\begin{aligned}\Delta(a) \bullet_\varepsilon \Delta(b) &= a^{(1)} \bullet b^{(1)} \otimes a^{(2)} b^{(2)} + \varepsilon(b^{(1)}) a^{(1)} \otimes a^{(2)} \bullet b^{(2)} \\ &= a^{(1)} \bullet b^{(1)} \otimes a^{(2)} b^{(2)} + a^{(1)} \otimes a^{(2)} \bullet b \\ &= \Delta(a \bullet b).\end{aligned}$$

Lemma 22. 1. Let A, B, C be three Com-PreLie algebras, $\varepsilon_A : A \rightarrow \mathbb{K}$ and $\varepsilon_B : B \rightarrow \mathbb{K}$ with the condition of lemma 21. Then $\varepsilon_A \otimes \varepsilon_B : A \otimes B \rightarrow \mathbb{K}$ also satisfies the condition of lemma 21. Moreover, the Com-PreLie algebras $(A \otimes B) \otimes C$ and $A \otimes (B \otimes C)$ are equal.

2. Let A, B be two Com-PreLie algebras, and $\varepsilon : A \rightarrow \mathbb{K}$ such that:

$$\forall a, b \in A, \quad \varepsilon(ab) = \varepsilon(a)\varepsilon(b), \quad \varepsilon(a \bullet b) = 0.$$

Then $\varepsilon \otimes Id : A \otimes B \rightarrow B$ is morphism of Com-PreLie algebras.

3. Let A, A', B, B' be Com-PreLie algebras, $\varepsilon : A \rightarrow \mathbb{K}$ and $\varepsilon' : A' \rightarrow \mathbb{K}$ satisfying the condition of lemma 21. Let $f : A \rightarrow A'$, $g : B \rightarrow B'$ be Com-PreLie algebra morphisms such that $\varepsilon' \circ f = \varepsilon$. Then $f \otimes g : A \otimes B \rightarrow A' \otimes B'$ is a Com-PreLie algebra morphism.

Proof. 1. Indeed, if $a_1, a_2 \in A$, $b_1, b_2 \in B$:

$$\begin{aligned}\varepsilon_A \otimes \varepsilon_B((a_1 \otimes b_1) \bullet (a_2 \otimes b_2)) &= \varepsilon_A(a_1 \bullet a_2)\varepsilon_B(b_1 b_2) + \varepsilon_A(a_1)\varepsilon_A(a_2)\varepsilon_B(b_1 \bullet b_2) \\ &= \varepsilon_A(a_2 \bullet a_1)\varepsilon_B(b_2 b_1) + \varepsilon_A(a_2)\varepsilon_A(a_1)\varepsilon_B(b_2 \bullet b_1) \\ &= \varepsilon_A \otimes \varepsilon_B((a_2 \otimes b_2) \bullet (a_1 \otimes b_1)).\end{aligned}$$

Let $a_1, a_2 \in A$, $b_1, b_2 \in B$, $c_1, c_2 \in C$. In $(A \otimes B) \otimes C$:

$$\begin{aligned}(a_1 \otimes b_1 \otimes c_1) \bullet (a_2 \otimes b_2 \otimes c_2) &= ((a_1 \otimes b_1) \bullet (a_2 \otimes b_2)) \otimes c_1 c_2 + \varepsilon_A \otimes \varepsilon_B(a_2 \otimes b_2) a_1 \otimes b_1 \otimes c_1 \bullet c_2 \\ &= a_1 \bullet a_2 \otimes b_1 b_2 \otimes c_1 c_2 + \varepsilon_A(a_2) a_1 \otimes b_1 \bullet b_2 \otimes c_1 c_2 + \varepsilon_A(a_2)\varepsilon_B(b_2) a_1 \otimes b_1 \otimes c_1 \bullet c_2.\end{aligned}$$

In $A \otimes (B \otimes C)$:

$$\begin{aligned}(a_1 \otimes b_1 \otimes c_1) \bullet (a_2 \otimes b_2 \otimes c_2) &= a_1 \bullet a_2 \otimes b_1 b_2 \otimes c_1 c_2 + \varepsilon_A(a_2) a_1 \otimes ((b_1 \otimes c_1) \bullet (b_2 \otimes c_2)) \\ &= a_1 \bullet a_2 \otimes b_1 b_2 \otimes c_1 c_2 + \varepsilon_A(a_2) a_1 \otimes b_1 \bullet b_2 \otimes c_1 c_2 + \varepsilon_A(a_2)\varepsilon_B(b_2) a_1 \otimes b_1 \otimes c_1 \bullet c_2.\end{aligned}$$

So $(A \otimes B) \otimes C = A \otimes (B \otimes C)$.

2. Let $a_1, a_2 \in A$, $b_1, b_2 \in B$.

$$\begin{aligned}\varepsilon \otimes Id((a_1 \otimes b_1)(a_2 \otimes b_2)) &= \varepsilon \otimes Id((a_1 \otimes b_1) \bullet (a_2 \otimes b_2)) \\ &= \varepsilon(a_1 a_2) b_1 b_2 &= \varepsilon(a_1 \bullet a_2) b_1 b_2 + \varepsilon(a_1)\varepsilon(a_2) b_1 \bullet b_2 \\ &= \varepsilon(a_1)\varepsilon(a_2) b_1 b_2 &= \varepsilon(a_1)\varepsilon(a_2) b_1 \bullet b_2 \\ &= \varepsilon \otimes Id((a_1 \otimes b_1)\varepsilon \otimes Id(a_2 \otimes b_2)), &= \varepsilon \otimes Id((a_1 \otimes b_1) \bullet \varepsilon \otimes Id(a_2 \otimes b_2)).\end{aligned}$$

So $\varepsilon \otimes Id$ is a morphism.

3. $f \otimes g$ is obviously an algebra morphism. If $a_1, a_2 \in A$, $b_1, b_2 \in B$:

$$\begin{aligned}(f \otimes g)((a_1 \otimes b_1) \bullet (a_2 \otimes b_2)) &= (f \otimes g)(a_1 \bullet a_2 \otimes b_1 b_2 + \varepsilon(a_2) a_1 \otimes b_1 \bullet b_2) \\ &= f(a_1) \bullet f(a_2) \otimes g(b_1)g(b_2) + \varepsilon(f(a_2))f(a_1) \otimes g(b_1) \bullet g(b_2) \\ &= (f(a_1) \otimes g(b_1)) \bullet (f(a_2) \otimes g(b_2)).\end{aligned}$$

So $f \otimes g$ is a Com-PreLie algebra morphism. □

Lemma 23. *Let A be an associative commutative bialgebra, and V a subspace of A which generates A . Let \bullet be a product on A such that:*

$$\forall a, b, c \in A, \quad (ab) \bullet c = (a \bullet c)b + a(b \bullet c).$$

Then A is a Com-PreLie bialgebra if, and only if, for all $x \in V$, $b, c \in A$:

$$\begin{aligned} (x \bullet b) \bullet c - x \bullet (b \bullet c) &= (x \bullet c) \bullet b - x \bullet (c \bullet b), \\ \Delta(x \bullet b) &= x^{(1)} \otimes x^{(2)} \bullet b + x^{(1)} \bullet b^{(1)} \otimes x^{(2)} b^{(2)}. \end{aligned}$$

Proof. \implies . Obvious. \longleftarrow . We consider:

$$B = \{a \in A \mid \forall b, c \in A, (a \bullet b) \bullet c - a \bullet (b \bullet c) = (a \bullet c) \bullet b - a \bullet (c \bullet b)\}.$$

Copying the proof of lemma 2-1, we obtain that $1 \bullet b = 0$ for all $b \in A$. This easily implies that $1 \in B$. By hypothesis, $V \subseteq B$. Let $a_1, a_2 \in B$. For all $b, c \in A$:

$$\begin{aligned} &((a_1 a_2) \bullet b) \bullet c - (a_1 a_2) \bullet (b \bullet c) \\ &= ((a_1 \bullet b) \bullet c) a_2 + (a_1 \bullet b)(a_2 \bullet c) + (a_1 \bullet c)(a_2 \bullet b) + a_1((a_2 \bullet b) \bullet c) \\ &\quad - (a_1 \bullet (b \bullet c)) a_2 - a_1(a_2 \bullet (b \bullet c)) \\ &= ((a_1 \bullet b) \bullet c - a_1 \bullet (b \bullet c)) a_2 + a_1((a_2 \bullet b) \bullet c - a_2 \bullet (b \bullet c)) \\ &\quad + (a_1 \bullet b)(a_2 \bullet c) + (a_1 \bullet c)(a_2 \bullet b). \end{aligned}$$

As $a_1, a_2 \in B$, this is symmetric in b, c , so $a_1 a_2 \in B$. Hence, B is a unitary subalgebra of A which contains V , so is equal to A : A is Com-PreLie. Let us now consider:

$$C = \{a \in A \mid \forall b \in A, \Delta(a \bullet b) = a^{(1)} \otimes a^{(2)} \bullet b + a^{(1)} \bullet b^{(1)} \otimes a^{(2)} b^{(2)}\}.$$

By hypothesis, $V \subseteq C$. Let $b \in B$.

$$\emptyset \otimes \emptyset \bullet b + \emptyset \bullet b^{(1)} \otimes 1b^{(2)} = 0 = \Delta(\emptyset \bullet b),$$

so $\emptyset \in C$. Let $a_1, a_2 \in C$. For all $b \in A$:

$$\begin{aligned} \Delta((a_1 a_2) \bullet b) &= \Delta((a_1 \bullet b) a_2 + a_1(a_2 \bullet b)) \\ &= a_1^{(1)} a_2^{(1)} \otimes (a_1^{(2)} \bullet b) a_2^{(2)} + (a_1^{(1)} \bullet b^{(1)}) a_2^{(1)} \otimes a_1^{(2)} b^{(2)} a_2^{(2)} \\ &\quad + a_1^{(1)} a_2^{(1)} \otimes a_1^{(2)} (a_2^{(2)} \bullet b) + a_1^{(1)} (a_2^{(1)} \bullet b^{(1)}) \otimes a_1^{(2)} a_2^{(2)} b^{(2)} \\ &= a_1^{(1)} a_2^{(1)} \otimes (a_1^{(2)} a_2^{(2)}) \bullet b + (a_1^{(1)} a_2^{(1)}) \bullet b^{(1)} \otimes a_1^{(2)} a_2^{(2)} b^{(2)} \\ &= (a_1 a_2)^{(1)} \otimes (a_1 a_2)^{(2)} \bullet b + (a_1 a_2)^{(1)} \bullet b^{(1)} \otimes (a_1 a_2)^{(2)} b^{(2)}. \end{aligned}$$

Hence, $a_1 a_2 \in C$, and C is a unitary subalgebra of A . As it contains V , $C = A$ and A is a Com-PreLie Hopf algebra. \square

4.2 Coproduct on $UCP(\mathcal{D})$

Definition 24. 1. *Let T be a partitioned tree and $I \subseteq V(T)$. We shall say that I is an ideal of T if for any vertex $v \in I$ and any vertex $w \in V(T)$ such that there exists an edge from v to w , then $w \in I$. The set of ideals of T is denoted $\mathcal{Id}(T)$.*

2. *Let T be partitioned forest decorated by $\mathbb{N} \times I$, and $I \in \mathcal{Id}(T)$.*

- *By restriction, I is a partitioned decorated forest. The product \cdot of the trees of I is denoted by $P^I(F)$.*

- By restriction, $T \setminus I$ is a partitioned decorated tree. For any vertex $v \in T \setminus I$, if we denote by (i, d) the decoration of v in T , we replace it by $(i + \iota_I(v), d)$, where $\iota_I(v)$ is the number of blocks C of T , included in I , such that there exists an edge from v to any vertex of C . The partitioned decorated tree obtained in this way is denoted by $R^I(F)$.

Theorem 25. We define a coproduct on $UCP(\mathcal{D})$ in the following way:

$$\forall T \in \mathcal{PT}(\mathbb{N} \times \mathcal{D}), \quad \Delta(T) = \sum_{I \in \mathcal{Id}(T)} R^I(T) \otimes P^I(T).$$

Then $UCP(\mathcal{D})$ is a Com-PreLie bialgebra. Moreover, $CP(\mathcal{D})$ and $\mathcal{H}_{CK}^{\mathcal{D}}$ are Com-PreLie bialgebra quotients of $UCP(\mathcal{D})$, and $\mathcal{H}_{CK}^{\mathcal{D}}$ is the Connes-Kreimer Hopf algebra of decorated rooted trees [3, 7].

Proof. We consider:

$$\varepsilon : \begin{cases} UCP(\mathcal{D}) & \longrightarrow \mathbb{K} \\ F & \longrightarrow \delta_{F,1}. \end{cases}$$

By lemma 22-1, $UCP(\mathcal{D}) \otimes_{\varepsilon} UCP(\mathcal{D})$ is a Com-PreLie algebra. It is unitary, the unit being $1 \otimes 1$. Hence, there exists a unique Com-PreLie algebra morphism $\Delta' : UCP(\mathcal{D}) \longrightarrow UCP(\mathcal{D}) \otimes_{\varepsilon} UCP(\mathcal{D})$, sending $\cdot_{(0,d)}$ over $\cdot_{(0,d)} \otimes 1 + 1 \otimes \cdot_{(0,d)}$ for all $d \in \mathcal{D}$. By lemma 22-2, $(UCP(\mathcal{D}) \otimes_{\varepsilon} UCP(\mathcal{D})) \otimes_{\varepsilon \otimes \varepsilon} UPC(\mathcal{D})$ and $UCP(\mathcal{D}) \otimes_{\varepsilon} (UCP(\mathcal{D}) \otimes_{\varepsilon} UCP(\mathcal{D}))$ are equal, and as both $(Id \otimes \Delta') \circ \Delta'$ and $(\Delta' \otimes Id) \circ \Delta'$ are Com-PreLie algebra morphisms sending $\cdot_{(0,d)}$ over $\cdot_{(0,d)} \otimes 1 \otimes 1 + 1 \otimes \cdot_{(0,d)} \otimes 1 + 1 \otimes 1 \otimes \cdot_{(0,d)}$ for all $d \in \mathcal{D}$, they are equal: Δ' is coassociative. Moreover, $(Id \otimes \varepsilon) \circ \Delta'$ and $(\varepsilon \otimes Id) \circ \Delta'$ are Com-PreLie endomorphisms of $UCP(\mathcal{D})$ sending $\cdot_{(0,d)}$ over itself for all $d \in \mathcal{D}$, so they are both equal to Id : ε is the counit of Δ' . Hence, with this coproduct Δ' , $UCP(\mathcal{D})$ is a Com-PreLie bialgebra.

Let us now prove that $\Delta(T) = \Delta'(T)$ for all $T \in \mathcal{PT}(\mathbb{N} \times \mathcal{D})$. We proceed by induction on the number of vertices n of T . If $n = 0$ or $n = 1$, it is obvious. Let us assume the result at all ranks $< n$. If T has strictly more than one root, we can write $T = T' \cdot T''$, where T' and T'' has strictly less than n vertices. It is easy to see that the ideals of T are the parts of $T' \sqcup T''$ of the form $I' \sqcup I''$, such that $I' \in \mathcal{Id}(T')$ and $I'' \in \mathcal{Id}(T'')$. Moreover, for such an ideal of T ,

$$R^{I' \sqcup I''}(T' \cdot T'') = R^{I'}(T') \cdot R^{I''}(T''), \quad P^{I' \sqcup I''}(T' \cdot T'') = P^{I'}(T') \cdot P^{I''}(T'').$$

Hence:

$$\begin{aligned} \Delta(T) &= \sum_{I' \in \mathcal{Id}(T'), I'' \in \mathcal{Id}(T'')} R^{I'}(T') \cdot R^{I''}(T'') \otimes R^{I'}(T') R^{I''}(T'') \\ &= \Delta(T) \cdot \Delta(T'') \\ &= \Delta'(T') \cdot \Delta'(T'') \\ &= \Delta'(T \cdot T'') \\ &= \Delta(T). \end{aligned}$$

If T has only one root, we can write $T = \cdot_{(i,d)} \bullet (T_1 \times \dots \times T_k)$, where $T_1, \dots, T_k \in \mathcal{PT}(\mathbb{N} \times \mathcal{D})$. The induction hypothesis holds for T_1, \dots, T_k . The ideals of T are:

- T itself: for this ideal I , $P^I(T) = T$ and $R^I(T) = \emptyset$.
- Ideals $I_1 \sqcup \dots \sqcup I_k$, where I_j is an ideal of T_j for all j . For such an ideal I , $P^I(T) = P^{I_1}(T_1) \cdot \dots \cdot P^{I_k}(T_k)$. Let $J = \{i_1, \dots, i_p\}$ be the set of indices i such that $I_i = T_i$, that is

to say the number of blocks C of I such that is an edge from the root of T to any vertex of C . Then:

$$\begin{aligned}
R^I(T) &= \bullet_{(i+p, d)} \bullet \prod_{j \notin J}^{\times} R^{I_j}(T_j) \\
&= f_{UCP(\mathcal{D})}^I(\bullet_{(i, d)}) \bullet \prod_{j \notin J}^{\times} R^{I_j}(T_j) \\
&= \bullet_{(i, d)} \bullet \emptyset^{\times p} \times t \prod_{j \notin J}^{\times} R^{I_j}(T_j) \\
&= \bullet_{(i, d)} \bullet R^{I_1}(T_1) \times \dots \times R^{I_k}(T_k).
\end{aligned}$$

We used lemma 5 for the third equality.

By proposition 4, with $a = \bullet_{(i, d)}$ and $b_1 \times \dots \times b_n = T_1 \times \dots \times T_k$:

$$\begin{aligned}
\Delta'(T) &= \sum_{I \subseteq [k]} \bullet_{(i, d)} \bullet \left(\prod_{i \in I}^{\times} T_i^{(1)} \right) \otimes \left(\prod_{i \in I} T_i^{(2)} \right) \emptyset \bullet \left(\prod_{i \notin I}^{\times} T_i \right) \\
&+ \sum_{I \subseteq [k]} \emptyset \bullet \left(\prod_{i \in I}^{\times} T_i^{(1)} \right) \otimes \left(\prod_{i \in I} T_i^{(2)} \right) \bullet_{(i, d)} \bullet \left(\prod_{i \notin I}^{\times} T_i \right) \\
&= \bullet_{(i, d)} \bullet T_1^{(1)} \times \dots \times T_k^{(1)} \otimes T_1^{(2)} \cdot \dots \cdot T_k^{(2)} + 0 \\
&+ \emptyset \otimes \bullet_{(i, d)} \bullet T_1 \times \dots \times T_k \\
&= \sum_{I_j \in Id(T_j)} \bullet_{(i, d)} \bullet R^{I_1}(T_1) \times \dots \times R^{I_k}(T_k) \otimes P^{I_1}(T_1) \cdot \dots \cdot P^{I_k}(T_k) + \emptyset \otimes T \\
&= \sum_{I \in Id(T), I \neq T} R^I(T) \otimes P^I(T) + \emptyset \otimes T \\
&= \sum_{I \in Id(T)} R^I(T) \otimes P^I(T) \\
&= \Delta(T).
\end{aligned}$$

Hence, $\Delta' = \Delta$.

For all $d \in \mathcal{D}$, $\bullet_{(0, d)} - \bullet_{(1, d)}$ is primitive, so $\Delta(\bullet_{(0, d)} - \bullet_{(1, d)}) \in I \otimes UCP(\mathcal{D}) + UCP(\mathcal{D}) \otimes I$. Consequently, I is a coideal, and the quotient $UCP(\mathcal{D})/I = CP(\mathcal{D})$ is a Com-PreLie bialgebra.

Let $x, y \in CP(\mathcal{D})$. By proposition 4, as \bullet_d is primitive:

$$\Delta(\bullet_d \bullet (x \times y)) = \bullet_d \bullet (x^{(1)} \times y^{(1)}) \otimes x^{(2)} \cdot y^{(2)} + 1 \otimes \bullet_d \bullet (x \times y),$$

whereas, by the 1-cocycle property:

$$\Delta(\bullet_d \bullet (x \cdot y)) = \bullet_d \bullet (x^{(1)} \cdot y^{(1)}) \otimes x^{(2)} \cdot y^{(2)} + \otimes \bullet_d \bullet (x \cdot y).$$

Hence:

$$\begin{aligned}
\Delta(\bullet_d \bullet (x \times y) - \bullet_d \bullet (x \cdot y)) &= \underbrace{(\bullet_d \bullet (x^{(1)} \times y^{(1)}) - \bullet_d \bullet (x^{(1)} \cdot y^{(1)}))}_{\in J} \otimes x^{(2)} \cdot y^{(2)} \\
&+ 1 \otimes \underbrace{(\bullet_d \bullet (x \times y) - \bullet_d \bullet (x \cdot y))}_{\in J} \\
&\in J \otimes CP(\mathcal{D}) + CP(\mathcal{D}) \otimes J,
\end{aligned}$$

so J is a coideal and $CP(\mathcal{D})/J = \mathcal{H}_{CK}^{\mathcal{D}}$ is a Com-PreLie bialgebra.

Let us consider:

$$B_d : \begin{cases} \mathcal{H}_{CK}^{\mathcal{D}} & \longrightarrow \mathcal{H}_{CK}^{\mathcal{D}} \\ T_1 \dots T_k & \longrightarrow \bullet_d \bullet T_1 \times \dots \times T_k, \end{cases}$$

where T_1, \dots, T_k are rooted trees decorated by \mathcal{D} . In other terms, $B_d(T_1 \dots T_k)$ is the tree obtained by grafting the forest $T_1 \dots T_k$ on a common root decorated by d . By proposition 4 and lemma 5, for all forest $F = T_1 \dots T_k \in \mathcal{H}_{CK}^{\mathcal{D}}$:

$$\begin{aligned} \Delta \circ B_d(F) &= \bullet_d \bullet T_1^{(1)} \times \dots \times T_k^{(1)} \otimes T_1^{(2)} \dots T_k^{(2)} + 0 + \emptyset \otimes \bullet_d \bullet T_1 \times \dots \times T_k \\ &= B_d(F^{(1)}) \otimes F^{(2)} + \emptyset \otimes B_d(F). \end{aligned}$$

We recognize the 1-cocycle property which characterizes the Connes-Kreimer coproduct of rooted trees, so $\mathcal{H}_{CK}^{\mathcal{D}}$ is indeed the Connes-Kreimer Hopf algebra. \square

Example 6. Let $i, j, k \in \mathbb{N}$ and $d, e, f \in \mathcal{D}$. In $UCP(\mathcal{D})$:

$$\begin{aligned} \Delta \bullet_{(i,d)} &= \bullet_{(i,d)} \otimes \emptyset + \emptyset \otimes \bullet_{(i,d)}, \\ \Delta \mathbf{!}_{\binom{j,e}{i,d}} &= \mathbf{!}_{\binom{j,e}{i,d}} \otimes \emptyset + \emptyset \otimes \mathbf{!}_{\binom{j,e}{i,d}} + \bullet_{(i+1,d)} \otimes \bullet_{(j,e)}, \\ \Delta^{(j,e)} \mathbf{V}_{\binom{k,f}{i,d}} &= \binom{j,e}{i,d} \mathbf{V}_{\binom{k,f}{i,d}} \otimes \emptyset + \emptyset \otimes \binom{j,e}{i,d} \mathbf{V}_{\binom{k,f}{i,d}} \\ &\quad + \mathbf{!}_{\binom{j,e}{i+1,d}} \otimes \bullet_{(k,f)} + \mathbf{!}_{\binom{k,f}{i+1,d}} \otimes \bullet_{(j,e)} + \bullet_{(i+2,d)} \otimes \binom{j,e}{i,d} \mathbf{!}_{(k,f)}, \\ \Delta^{(j,e)} \mathbf{V}_{\binom{k,f}{i,d}} &= \binom{j,e}{i,d} \mathbf{V}_{\binom{k,f}{i,d}} \otimes \emptyset + \emptyset \otimes \binom{j,e}{i,d} \mathbf{V}_{\binom{k,f}{i,d}} \\ &\quad + \mathbf{!}_{\binom{j,e}{i,d}} \otimes \bullet_{(k,f)} + \mathbf{!}_{\binom{k,f}{i,d}} \otimes \bullet_{(j,e)} + \bullet_{(i+1,d)} \otimes \binom{j,e}{i,d} \mathbf{!}_{(k,f)}, \\ \Delta \mathbf{!}_{\binom{k,f}{i,d}} &= \mathbf{!}_{\binom{k,f}{i,d}} \otimes \emptyset + \emptyset \otimes \mathbf{!}_{\binom{k,f}{i,d}} + \mathbf{!}_{\binom{j,d+1,e}{i,d}} \otimes \bullet_{(k,f)} + \bullet_{(i+1,d)} \otimes \mathbf{!}_{\binom{k,f}{j,e}}. \end{aligned}$$

In $CP(\mathcal{D})$:

$$\begin{aligned} \Delta \bullet_d &= \bullet_d \otimes \emptyset + \emptyset \otimes \bullet_d, \\ \Delta \mathbf{!}_d^e &= \mathbf{!}_d^e \otimes \emptyset + \emptyset \otimes \mathbf{!}_d^e + \bullet_d \otimes \bullet_e, \\ \Delta^e \mathbf{V}_d^f &= {}^e \mathbf{V}_d^f \otimes \emptyset + \emptyset \otimes {}^e \mathbf{V}_d^f + \mathbf{!}_d^e \otimes \bullet_f + \mathbf{!}_d^f \otimes \bullet_e + \bullet_d \otimes e \mathbf{!}_{f}, \\ \Delta^e \mathbf{V}_d^f &= {}^e \mathbf{V}_d^f \otimes \emptyset + \emptyset \otimes {}^e \mathbf{V}_d^f + \mathbf{!}_d^e \otimes \bullet_f + \mathbf{!}_d^f \otimes \bullet_e + \bullet_d \otimes e \mathbf{!}_{f}, \\ \Delta \mathbf{!}_d^f &= \mathbf{!}_d^f \otimes \emptyset + \emptyset \otimes \mathbf{!}_d^f + \mathbf{!}_d^e \otimes \bullet_f + \bullet_d \otimes \mathbf{!}_e^f. \end{aligned}$$

In $\mathcal{H}_{CK}^{\mathcal{D}}$:

$$\begin{aligned} \Delta \bullet_d &= \bullet_d \otimes \emptyset + \emptyset \otimes \bullet_d, \\ \Delta \mathbf{!}_d^e &= \mathbf{!}_d^e \otimes \emptyset + \emptyset \otimes \mathbf{!}_d^e + \bullet_d \otimes \bullet_e, \\ \Delta^e \mathbf{V}_d^f &= {}^e \mathbf{V}_d^f \otimes \emptyset + \emptyset \otimes {}^e \mathbf{V}_d^f + \mathbf{!}_d^e \otimes \bullet_f + \mathbf{!}_d^f \otimes \bullet_e + \bullet_d \otimes \bullet_e \mathbf{!}_f, \\ \Delta \mathbf{!}_d^f &= \mathbf{!}_d^f \otimes \emptyset + \emptyset \otimes \mathbf{!}_d^f + \mathbf{!}_d^e \otimes \bullet_f + \bullet_d \otimes \mathbf{!}_e^f. \end{aligned}$$

4.3 An application: Connes-Moscovici subalgebras

Let us fix a set \mathcal{D} of decorations. For any $d \in \mathcal{D}$, we define an operator $N_d : \mathcal{H}_{CK}^{\mathcal{D}} \longrightarrow \mathcal{H}_{CK}^{\mathcal{D}}$ by:

$$\forall x \in \mathcal{H}_{CK}^{\mathcal{D}}, \quad N_d(x) = x \bullet \bullet_d.$$

In other words, if F is a rooted forest, $N_d(F)$ is the sum of all forests obtained by grafting a leaf decorated by d on a vertex of F : when \mathcal{D} is reduced to a singleton, this is the growth operator N of [3].

For all $k \geq 1$, $i_1, \dots, i_k \in \mathcal{D}$, we put:

$$X_{i_1, \dots, i_k} = N_{i_k} \circ \dots \circ N_{i_2}(\bullet_{i_1}).$$

When $|\mathcal{D}| = 1$, these are the generators of the Connes-Moscovici subalgebra of [3].

Proposition 26. Let $\mathcal{H}_{CM}^{\mathcal{D}}$ be the subalgebra of $\mathcal{H}_{CK}^{\mathcal{D}}$ generated by all the elements X_{i_1, \dots, i_k} . Then $\mathcal{H}_{CM}^{\mathcal{D}}$ is a Hopf subalgebra.

Proof. Note that N_d is a derivation; as $N_d(X_{i_1, \dots, i_k}) = X_{i_1, \dots, i_k, d}$ for all $i_1, \dots, i_k, d \in \mathcal{D}$, $\mathcal{H}_{CM}^{\mathcal{D}}$ is stable under N_d for any $d \in \mathcal{D}$. As the X_{i_1, \dots, i_k} are homogenous of degree k :

$$X_{i_1, \dots, i_k} \bullet 1 = kX_{i_1, \dots, i_k}.$$

Hence, $\mathcal{H}_{CM}^{\mathcal{D}}$ is stable under the derivation $D : x \mapsto x \bullet 1$. We obtain:

$$\begin{aligned} \Delta(X_{i_1, \dots, i_k}) &= \Delta(X_{i_1, \dots, i_{k-1}} \bullet \cdot i_k) \\ &= X_{i_1, \dots, i_{k-1}}^{(1)} \otimes X_{i_1, \dots, i_{k-1}}^{(2)} \bullet \cdot i_k \\ &\quad + X_{i_1, \dots, i_{k-1}}^{(1)} \bullet \cdot i_k \otimes X_{i_1, \dots, i_{k-1}}^{(2)} + X_{i_1, \dots, i_{k-1}}^{(1)} \bullet \emptyset \otimes X_{i_1, \dots, i_{k-1}}^{(2)} \bullet \cdot i_k. \end{aligned} \tag{10}$$

An easy induction on k proves that $\Delta(X_{i_1, \dots, i_k})$ belongs to $\mathcal{H}_{CM}^{\mathcal{D}} \otimes \mathcal{H}_{CM}^{\mathcal{D}}$. \square

Proposition 27. We assume that \mathcal{D} is finite. Then $\mathcal{H}_{CM}^{\mathcal{D}}$ is the graded dual of the enveloping algebra of the augmentation ideal of the Com-PreLie algebra $T(V, f)$, where $V = \text{Vect}(\mathcal{D})$ and $f = \text{Id}_V$.

Proof. We put $W = \text{Vect}(X_{i_1, \dots, i_k} \mid k \geq 1, i_1, \dots, i_k \in \mathcal{D})$. As this is the case for $\mathcal{H}_{CK}^{\mathcal{D}}$, for any $x \in W$:

$$\Delta(x) - x \otimes 1 + 1 \otimes x \in W \otimes \mathcal{H}_{CM}^{\mathcal{D}}.$$

This implies that the graded dual of $\mathcal{H}_{CM}^{\mathcal{D}}$ is the enveloping of a graded algebra \mathfrak{g} ; as a vector space, \mathfrak{g} is identified with W^* and its preLie product is dual of the bracket δ defined on W by $(\pi_W \otimes \pi_W \circ \Delta)$, where π_W is the canonical projection on W which vanishes on $(1) + (\mathcal{H}_{CM}^{\mathcal{D}})_+^2$. By (10), using Sweedler's notation $\delta(x) = x' \otimes x''$, we obtain:

$$\delta(X_{i_1, \dots, i_{k+1}}) = X'_{i_1, \dots, i_k} \otimes X''_{i_1, \dots, i_k} \bullet X_{i_{k+1}} + X'_{i_1, \dots, i_k} \bullet X_{i_{k+1}} \otimes X''_{i_1, \dots, i_k} + kX_{i_1, \dots, i_k} \otimes X_{i_{k+1}}.$$

We shall use the following notations. If $I \subseteq [k]$, we put:

- $m(I) = \max(i \mid [i] \subseteq I)$, with the convention $m(I) = 0$ if $1 \notin I$.
- $X_{i_I} = X_{i_{p_1}, \dots, i_{p_l}}$ if $I = \{p_1 < \dots < p_l\}$.

An easy induction then proves the following result:

$$\forall i_1, \dots, i_k \in \mathcal{D}, \quad \delta(X_{i_1, \dots, i_k}) = \sum_{\emptyset \subsetneq I \subseteq [k]} m(I) X_{i_I} \otimes X_{i_{[k] \setminus I}}.$$

We identify W^* and $T(V)_+$ via the pairing:

$$\forall i_1, \dots, i_k, j_1, \dots, j_l \in \mathcal{D}, \quad \langle X_{i_1, \dots, i_k}, j_1 \dots j_l \rangle = \delta_{(i_1, \dots, i_k), (j_1, \dots, j_l)}.$$

The preLie product on $T(V)_+$ induced by δ is then given by:

$$i_1 \dots i_k \bullet i_{k+1} \dots i_{k+l} = \sum_{\sigma \in \text{Sh}(k, l)} m_k(\sigma) i_{\sigma^{-1}(1)} \dots i_{\sigma^{-1}(k+l)}.$$

By (7), this is precisely the preLie product of $T(V, f)$. \square

Remark 9. The following map is a bijection:

$$\theta_{k,l} : \begin{cases} Sh(k,l) & \longrightarrow & Sh(l,k) \\ \sigma & \longrightarrow & (k+l \ k+l-1 \dots 1) \circ \sigma \circ (k+l \ k+l-1 \dots 1). \end{cases}$$

Moreover, for any $\sigma \in Sh(k,l)$:

$$m_l(\theta_{k,l}(\sigma)) = \min\{i \in l \in \{k+1, \dots, k+l\} \mid \sigma(i) = i, \dots, \sigma(k+l) = \sigma(k+l)\} = m'_l(\sigma),$$

with the convention $m'_l(\sigma) = 0$ if $\sigma(k+l) \neq k+l$. Then the Lie bracket associated to \bullet is given by:

$$\forall i_1, \dots, i_{k+l} \in \mathcal{D}, \quad [i_1 \dots i_k, i_{k+1} \dots i_{k+l}] = \sum_{\sigma \in Sh(k,l)} (m_k(\sigma) - m'_l(\sigma)) i_{\sigma^{-1}(1)} \dots i_{\sigma^{-1}(k+l)}.$$

4.4 A rigidity theorem for Com-PreLie bialgebras

Theorem 28. *Let (A, m, \bullet, Δ) be a connected Com-PreLie bialgebra. If f_A (defined in Proposition 3) is surjective, then (A, m, Δ) and $(T(\text{Prim}(A)), \sqcup, \Delta)$ are isomorphic Hopf algebras.*

Proof. We put $V = \text{Prim}(A)$.

First step. As f_A is surjective, there exists $g : V \longrightarrow V$ such that $f_A \circ g = \text{Id}_V$. For all $x \in V$, we put:

$$L_x : \begin{cases} A & \longrightarrow & A \\ y & \longrightarrow & g(x) \bullet y. \end{cases}$$

For all $y \in A$:

$$\Delta \circ L_x(y) = \emptyset \otimes g(x) \bullet y + g(x) \bullet y^{(1)} \otimes y^{(2)} = \emptyset \otimes L_x(y) + (\text{Id} \otimes L_x) \circ \Delta(y).$$

Hence, L_x is a 1-cocycle of A . Moreover, $L_x(1) = g(x) \bullet 1 = f_A \circ g(x) = x$. For all $x_1, \dots, x_n \in V$, we define $\omega(x_1, \dots, x_n)$ inductively on n by:

$$\omega(x_1, \dots, x_n) = \begin{cases} \emptyset & \text{if } n = 0, \\ L_{x_1}(\omega(x_2, \dots, x_{n-1})) & \text{if } n \geq 1. \end{cases}$$

In particular, $\omega(v) = v$ for all $v \in V$. An easy induction proves that:

$$\Delta(\omega(x_1, \dots, x_n)) = \sum_{i=0}^n \omega(x_1, \dots, x_i) \otimes \omega(x_{i+1}, \dots, x_n).$$

Hence, the following map is a coalgebra morphism:

$$\omega : \begin{cases} T(V) & \longrightarrow & A \\ x_1 \dots x_n & \longrightarrow & \omega(x_1, \dots, x_n). \end{cases}$$

It is injective: if $\text{Ker}(\omega)$ is nonzero, then it is a nonzero coideal of $T(V)$, so it contains nonzero primitive elements of $T(V)$, that is to say nonzero elements of V . For all $v \in V$, $\omega(v) = L_v(1) = v$: contradiction. Let us prove that ω is surjective. As A is connected, for any $x \in A_+$, there exists $n \geq 1$ such that $\tilde{\Delta}^{(n)}(x) = 0$. Let us prove that $x \in \text{Im}(\omega)$ by induction on n . If $n = 1$, then $x \in V$, so $x = \omega(x)$. Let us assume the result at all ranks $< n$. By coassociativity of $\tilde{\Delta}$, $\tilde{\Delta}^{(n-1)}(x) \in V^{\otimes n}$. We put $\tilde{\Delta}^{(n-1)}(x) = x_1 \otimes \dots \otimes x_n \in V^{\otimes n}$. Then $\tilde{\Delta}^{(n-1)}(x) = \tilde{\Delta}^{(n-1)}(\omega(x_1, \dots, x_n))$. By the induction hypothesis, $x - \omega(x_1, \dots, x_n) \in \text{Im}(\omega)$, so $x \in \text{Im}(\omega)$.

We proved that the coalgebras A and $T(V)$ are isomorphic. We now assume that $A = T(V)$ as a coalgebra.

Second step. We denote by π the canonical projection on V in $T(V)$. Let $\varpi : T_+(V) \longrightarrow V$ be any linear map. We define:

$$F_\varpi : \begin{cases} T(V) & \longrightarrow T(V) \\ x_1 \dots x_n & \longrightarrow \sum_{k=1}^n \sum_{i_1+\dots+i_k=n} \varpi(x_1 \dots x_{i_1}) \dots \varpi(x_{i_1+\dots+i_{k-1}+1} \dots x_n). \end{cases}$$

Let us prove that F_ϖ is the unique coalgebra endomorphism such that $\pi \circ F_\varpi = \varpi$. First:

$$\begin{aligned} \Delta(F_\varpi(x_1 \dots x_n)) &= \sum_{i_1+\dots+i_k=n} \Delta(\varpi(x_1 \dots x_{i_1}) \dots \varpi(x_{i_1+\dots+i_{k-1}+1} \dots x_n)) \\ &= \sum_{i_1+\dots+i_k=n} \sum_{j=0}^k \varpi(x_1 \dots x_{i_1}) \dots \varpi(x_{i_1+\dots+i_{j-1}+1} \dots x_{i_1+\dots+i_j}) \\ &\quad \otimes \varpi(x_{i_1+\dots+i_j+1} \dots x_{i_1+\dots+i_{j+1}}) \dots \varpi(x_{i_1+\dots+i_{k-1}+1} \dots x_n) \\ &= \sum_{i=0}^n F_\varpi(x_1 \dots x_i) \otimes F_\varpi(x_{i+1} \dots x_n) \\ &= (F_\varpi \otimes F_\varpi) \circ \Delta(x_1 \dots x_n). \end{aligned}$$

Moreover:

$$\begin{aligned} \pi \circ F_\varpi(x_1 \dots x_n) &= \sum_{k=1}^n \sum_{i_1+\dots+i_k=n} \pi(\varpi(x_1 \dots x_{i_1}) \dots \varpi(x_{i_1+\dots+i_{k-1}+1} \dots x_n)) \\ &= \pi \circ \varpi(x_1 \dots x_n) + 0 \\ &= \varpi(x_1 \dots x_n). \end{aligned}$$

Let us now prove the unicity. Let F, G be two coalgebra endomorphisms such that $\pi \circ F = \pi \circ G = \varpi$. If $F \neq G$, let $x_1 \dots x_n$ be a word of $T(V)$, such that $F(x_1 \dots x_n) - G(x_1 \dots x_n) \neq 0$, of minimal length. By minimality of n :

$$\tilde{\Delta}(F(x_1 \dots x_n)) = (F \otimes F) \circ \tilde{\Delta}(x_1 \dots x_n) = (G \otimes G) \circ \tilde{\Delta}(x_1 \dots x_n) = \tilde{\Delta}(G(x_1 \dots x_n)).$$

Hence, $F(x_1 \dots x_n) - G(x_1 \dots x_n) \in \text{Prim}(T(V)) = V$, so:

$$F(x_1 \dots x_n) - G(x_1 \dots x_n) = \pi(F(x_1 \dots x_n) - G(x_1 \dots x_n)) = \varpi(x_1 \dots x_n) - \varpi(x_1 \dots x_n) = 0.$$

This is a contradiction, so $F = G$.

Third step. Let $\varpi_1, \varpi_2 : T_+(V) \longrightarrow V$ and let $F_1 = F_{\varpi_1}$, $F_2 = F_{\varpi_2}$ be the associated coalgebra morphisms. Then:

$$\pi \circ F_2 \circ F_1(x_1 \dots x_n) = \sum_{i_1+\dots+i_k=n} \varpi_2(\varpi_1(x_1 \dots x_{i_1}) \dots \varpi_1(x_{i_1+\dots+i_{k-1}+1} \dots x_n)).$$

We denote this map by $\varpi_2 \diamond \varpi_1$. By the unicity in the second step, $F_2 \circ F_1 = F_{\varpi_2 \diamond \varpi_1}$. It is not difficult to prove that for any $\varpi : T_+(V) \longrightarrow V$, there exists $\varpi' : T_+(V) \longrightarrow V$, such that $\varpi' \diamond \varpi = \varpi \diamond \varpi' = \pi$ if, and only if, $\varpi|_V$ is invertible. If this holds, then $F_\varpi \circ F_{\varpi'} = F_{\varpi'} \circ F_\varpi = F_\pi = Id$, by the unicity in the second step. So, if $\varpi|_V$ is invertible, then F_ϖ is invertible.

Fourth step. We denote by $*$ the product of $T(V)$. Let us choose $\varpi : T_+(V) \rightarrow V$ such that $\varpi(T_+(V) * T_+(V)) = (0)$. Let $F = F_\varpi$ the associated coalgebra morphism. As \emptyset is the unique group-like element of $T(V)$, the unit of $*$ is \emptyset . Let us prove that for all $x, y \in T(V)$, $F(x * y) = F(x) \cdot F(y)$. We proceed by induction on $\text{length}(x) + \text{length}(y) = n$. As \emptyset is the unit for both $*$ and \cdot and $F(\emptyset) = \emptyset$, it is obvious if x or y is equal to \emptyset : this observation covers the case $n = 0$. Let us assume the result at all rank $< n$. By the preceding observation on the unit, we can assume that $x, y \in T_+(V)$. We put $G = F \circ *$ and $H = \cdot \circ (F \otimes F)$. They are both coalgebra morphisms from $T(V) \otimes T(V)$ to $T(V)$. Moreover:

$$\pi \circ G(x \otimes y) = \pi \circ F(x * y) = \varpi(x * y) = 0.$$

As the shuffle product is graded for the length, $\pi \circ H(x \otimes y) = 0$. By the induction hypothesis:

$$\tilde{\Delta} \circ G(x \otimes y) = (G \otimes G) \circ \tilde{\Delta}(x \otimes y) = (F \otimes F) \circ \tilde{\Delta}(x \otimes y) = \tilde{\Delta} \circ F(x \otimes y).$$

Hence, $G(x \otimes y) - F(x \otimes y)$ is primitive, so belongs to V . This implies:

$$G(x \otimes y) - F(x \otimes y) = \pi(G(x \otimes y) - F(x \otimes y)) = 0 - 0 = 0.$$

So $F(x * y) = G(x \otimes y) = F(x \otimes y) = F(x) \sqcup F(y)$. Hence, F is a bialgebra morphism from $(T(V), *, \Delta)$ to $(T(V), \sqcup, \Delta)$.

By the third and fourth steps, in order to prove that $(T(V), *, \Delta)$ and $(T(V), \sqcup, \Delta)$ are isomorphic, it is enough to find $\varpi : T_+(V) \rightarrow V$, such that $\varpi|_V$ is invertible and $\varpi(T_+(V) * T_+(V)) = (0)$; hence, it is enough to prove that $V \cap (A_+ * A_+) = (0)$.

Last step. We define $\Delta : \text{End}(A) \rightarrow \text{End}(A \otimes A, A)$ by $\Delta(f)(x \otimes y) = f(x * y)$. We denote by \star the convolution product of $\text{End}(A)$ induced by the bialgebra $(A, *, \Delta)$. Let $f, g \in \text{End}(A)$. We assume that we can write $\Delta(f) = f^{(1)} \otimes f^{(2)}$ and $\Delta(g) = g^{(1)} \otimes g^{(2)}$, that is to say, for all $x, y \in A$:

$$f(xy) = f^{(1)}(x) * f^{(2)}(y), \quad g(xy) = g^{(1)}(x) * g^{(2)}(y).$$

Then, as $*$ is commutative:

$$\begin{aligned} f \star g(x * y) &= f(x^{(1)} * y^{(1)}) * g(x^{(2)} * y^{(2)}) \\ &= f^{(1)}(x^{(1)}) * f^{(2)}(y^{(1)}) * g^{(1)}(x^{(2)}) * g^{(2)}(y^{(2)}) \\ &= f^{(1)}(x^{(1)}) * g^{(1)}(x^{(2)}) * f^{(2)}(y^{(1)}) * g^{(2)}(y^{(2)}) \\ &= f^{(1)} \star g^{(1)}(x) * f^{(2)} \star g^{(2)}(y). \end{aligned}$$

Hence, $\Delta(f \star g) = \Delta(f) \star \Delta(g)$.

Let ρ be the canonical projection on A_+ and 1 be the unit of the convolution algebra $\text{End}(V)$. Then $1 + \rho = \text{Id}$. As $\Delta(\text{Id}) = \text{Id} \otimes \text{Id}$ and $\Delta(1) = 1 \otimes 1$, this gives:

$$\Delta(\rho) = \rho \otimes 1 + 1 \otimes \rho + \rho \otimes \rho.$$

We consider:

$$\psi = \ln(1 + \rho) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \rho^{\star n}.$$

As A is connected, for all $x \in A$, $\rho^{\star n}(x) = 0$ if n is great enough, so ψ exists. Moreover, as Δ is compatible with the convolution product:

$$\begin{aligned} \Delta(\psi) &= \ln(1 \otimes 1 + \rho \otimes 1 + 1 \otimes \rho + \rho \otimes \rho) \\ &= \ln((1 + \rho) \otimes (1 + \rho)) \\ &= \ln(1 + \rho) \otimes 1 + \ln(1 \otimes (1 + \rho)) \\ &= \ln(1 + \rho) \otimes 1 + 1 \otimes \ln(1 + \rho) \\ &= \psi \otimes 1 + 1 \otimes \psi. \end{aligned}$$

We used $((1 + \rho) \otimes 1) \star (1 \otimes (1 + \rho)) = (1 \otimes (1 + \rho)) \star ((1 + \rho) \otimes 1) = (1 + \rho) \otimes (1 + \rho)$ for the third equality. Hence, for all $x, y \in A$:

$$\psi(x * y) = \psi(x)\varepsilon(y) + \varepsilon(x)\psi(y).$$

In particular, if $x, y \in A_+$, $\psi(x * y) = 0$. If $x \in V$, then $\rho^1(x) = x$ and if $n \geq 2$:

$$\rho^{*n}(x) = \sum_{i=1}^n \rho(1) * \dots * \rho(1) * \rho(x) * \rho(1) * \dots * \rho(1) = 0.$$

So $\psi(x) = x$. Finally, if $x \in V \cap (A_+ * A_+)$, $\psi(x) = x = 0$. So $V \cap (A_+ * A_+) = (0)$. \square

The following result is proved for $\mathcal{H}_{CK}^{\mathcal{D}}$ in [2] and in [7]:

Corollary 29. *$CP(\mathcal{D})$ and $\mathcal{H}_{CK}^{\mathcal{D}}$ are, as Hopf algebras, isomorphic to shuffle algebras.*

Proof. $CP(\mathcal{D})$ is a connected Com-PreLie bialgebra. Moreover, if $x \in CP(\mathcal{D})$, homogeneous of degree n , $x \bullet \emptyset = nx$. Hence, as the homogeneous component of degree 0 of $Prim(CP(\mathcal{D}))$ is zero, $f_{CP(\mathcal{D})}$ is invertible. By the rigidity theorem, $f_{CP(\mathcal{D})}$ is, as a Hopf algebra, isomorphic to a shuffle algebra. The proof is similar for $\mathcal{H}_{CK}^{\mathcal{D}}$. \square

Remark 10. 1. This is not the case for $UCP(\mathcal{D})$. For example, if d, e are two distinct elements of \mathcal{D} , it is not difficult to prove that there is no element $x \in UCK(\mathcal{D})$ such that:

$$\Delta(x) = x \otimes 1 + 1 \otimes x + \bullet_{(0,d)} \otimes \bullet_{(0,e)}.$$

So $UCP(\mathcal{D})$ is not cofree.

2. $CP(\mathcal{D})$ and $\mathcal{H}_{CK}^{\mathcal{D}}$ are not isomorphic, as Com-PreLie bialgebras, to any $T(V, f)$. Indeed, in $T(V, f)$, for any $x \in V$ such that $f(x) = x$, $x \sqcup x = 2x \bullet x = 2xx$. In $f_{CP(\mathcal{D})}$ or $\mathcal{H}_{CK}^{\mathcal{D}}$, for any $d \in \mathcal{D}$, with $x = \bullet_d$, $f(x) = x$ but $x \cdot x \neq 2x \bullet x$.

4.5 Dual of $UCP(\mathcal{D})$ and $CP(\mathcal{D})$

We identify $UCP(\mathcal{D})$ and its graded dual by considering the basis of partitioned trees as orthonormal; similarly, we identify $CP(\mathcal{D})$ and $\mathcal{H}_{CK}^{\mathcal{D}}$ with their graded dual.

Let us consider the Hopf algebra $(UCP(\mathcal{D}), \cdot, \Delta)$. As a commutative algebra, it is freely generated by the set $UPT_1(\mathcal{D})$ of partitioned trees decorated by $\mathbb{N} \times \mathcal{D}$ with one root. Moreover, if $T \in UPT_1(\mathcal{D})$:

$$\Delta(T) - 1 \otimes T \in Vect(UPT_1(\mathcal{D})) \otimes UCP(\mathcal{D}).$$

Consequently, this is a right-sided combinatorial bialgebra in the sense of [12], and its graded dual is the enveloping algebra of a PreLie algebra $\mathfrak{g}_{UCP(\mathcal{D})}$. Direct computations prove the following result:

Theorem 30. *The PreLie algebra $\mathfrak{g}_{UCP(\mathcal{D})}$ is the linear span of $UPT_1(\mathcal{D})$. For any $T, T' \in UPT_1(\mathcal{D})$, the PreLie product is given by:*

$$T \diamond T' = \sum_{\substack{s \in V(T), \\ b \in bl(s) \sqcup \{*\}}} (T \bullet_{s,b} T')[-1]_s.$$

Example 7. If $\mathcal{D} = \{1\}$, forgetting the second decoration of the vertices, in $\mathfrak{g}_{UCP(\mathcal{D})}$:

$$\begin{aligned} \bullet_i \diamond \bullet_j &= (1 - \delta_{i,0}) \mathfrak{!}_{i-1}^j, \\ \mathfrak{!}_i^j \diamond \bullet_k &= (1 - \delta_{j,0}) \mathfrak{!}_i^{k-1} + (1 - \delta_{i,0}) \left({}^j \mathfrak{V}_{i-1}^k + {}^j \mathfrak{V}_{i-1}^k \right). \end{aligned}$$

Similarly, the Hopf algebra $(CP(\mathcal{D}), \cdot, \Delta)$ is, as a commutative algebra, freely generated by the set $\mathcal{PT}_1(\mathcal{D})$ of partitioned trees decorated by \mathcal{D} with one root. Moreover, if $T \in \mathcal{PT}_1(\mathcal{D})$,

$$\Delta(T) - 1 \otimes T \in Vect(\mathcal{PT}_1(\mathcal{D})) \otimes CP(\mathcal{D}).$$

Consequently, its graded dual is the enveloping algebra of a PreLie algebra $\mathfrak{g}_{CP}(\mathcal{D})$, described by the following theorem:

Theorem 31. *The PreLie algebra $\mathfrak{g}_{CP}(\mathcal{D})$ is the linear span of $\mathcal{PT}_1(\mathcal{D})$. For any $T, T' \in \mathcal{PT}_1(\mathcal{D})$, the PreLie product is given by:*

$$T \diamond T' = \sum_{\substack{s \in V(T), \\ b \in bl(s) \sqcup \{*\}}} T \bullet_{s,b} T'.$$

Example 8. If $\mathcal{D} = \{1\}$, forgetting the decorations, in $\mathfrak{g}_{CP}(\mathcal{D})$:

$$\cdot \diamond \cdot = \mathfrak{!}, \quad \mathfrak{!} \diamond \cdot = \mathfrak{!} + \mathfrak{V} + \mathfrak{V}.$$

Notations 3. Let $T \in \mathcal{PT}_1(\mathcal{D})$. We can write $T = \cdot_a \bullet (T_1 \times \dots \times T_k) = B_d(T_1 \dots T_k)$, where $T_1, \dots, T_k \in \mathcal{PT}(\mathcal{D})$. Up to a change of indexation, we will always assume that $T_1, \dots, T_p \in \mathcal{PT}_1(\mathcal{D})$ and $T_{p+1}, \dots, T_k \notin \mathcal{PT}_1(\mathcal{D})$. The integer p is denoted by $\varsigma(T)$.

Proposition 32. *As a PreLie algebra, $\mathfrak{g}_{CP}(\mathcal{D})$ is freely generated by the set of trees $T \in \mathcal{PT}_1(\mathcal{D})$ such that $\varsigma(T) = 0$.*

Proof. We define a coproduct on $\mathfrak{g}_{CP}(\mathcal{D})$ in the following way:

$$\forall T = B_d(T_1 \dots T_k) \in \mathcal{PT}_1(\mathcal{D}), \quad \delta(T) = \sum_{i=1}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i.$$

This coproduct is permutative: indeed,

$$(\delta \otimes Id) \circ \delta(T) = \sum_{1 \leq i \neq j \leq \varsigma(T)} B_d(T_1 \dots \widehat{T}_i \dots \widehat{T}_j \dots T_k) \otimes T_i \otimes T_j,$$

so $(\delta \otimes Id) \circ \delta = (23) \cdot (\delta \otimes Id) \circ \delta$. Let $T = B_d(T_1 \dots T_k), T' \in \mathcal{PT}_1(\mathcal{D})$. Then:

$$T \diamond T' = B_d(T' T_1 \dots T_k) + \sum_{i=1}^k B_d(T_1 \dots (T_i \diamond T') \dots T_k) + \sum_{i=1}^k B_d(T_1 \dots (T_i \sqcup T') \dots T_k).$$

Hence:

$$\begin{aligned}
\delta(T \otimes T') &= B_d(T_1 \dots T_k) \otimes T' + \sum_{i=1}^{\varsigma(T)} B_d(T' T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \\
&+ \sum_{i=1}^k \sum_{\substack{j=1 \\ j \neq i}}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_j \dots (T_i \diamond T') \dots T_k) \otimes T_j + \sum_{i=1}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \diamond T' \\
&+ \sum_{i=1}^k \sum_{\substack{j=1 \\ j \neq i}}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_j \dots (T_i \sqcup T') \dots T_k) \otimes T_j \\
&= \sum_{j=1}^{\varsigma(T)} \left(B_d(T' T_1 \dots \widehat{T}_j \dots T_k) + \sum_{\substack{i=1 \\ i \neq j}}^k B_d(T_1 \dots \widehat{T}_j \dots (T_i \diamond T' + T_i \sqcup T') \dots T_k) \right) \otimes T_j \\
&+ \sum_{i=1}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \diamond T' + T \otimes T' \\
&= \sum_{j=1}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_j \dots T_k) \bullet T' \otimes T_j + \sum_{i=1}^{\varsigma(T)} B_d(T_1 \dots \widehat{T}_i \dots T_k) \otimes T_i \diamond T' + T \otimes T' \\
&= T^{(1)} \diamond T' \otimes T^{(2)} + T^{(1)} \otimes T^{(2)} \diamond T' + T \otimes T'.
\end{aligned}$$

By Livernet's rigidity theorem, $\mathfrak{g}_{CP}(\mathcal{D})$ is freely generated, as a PreLie algebra, by $Ker(\delta)$.

We define:

$$\Upsilon : \begin{cases} \mathfrak{g}_{CP}(\mathcal{D}) \otimes \mathfrak{g}_{CP}(\mathcal{D}) & \longrightarrow \mathfrak{g}_{CP}(\mathcal{D}) \\ T \otimes T' & \longrightarrow T \bullet_{r(T),*} T', \end{cases}$$

where $r(T)$ is the root of T . In other words, $\Upsilon(B_d(T_1 \dots T_k) \otimes T') = B_d(T' T_1 \dots T_k)$; this implies that for any $T \in \mathcal{PT}_1(\mathcal{D})$, $\Upsilon \circ \delta(T) = \varsigma(T)T$. Hence, if $x = \sum a_T T \in Ker(\delta)$, $\Upsilon \circ \delta(x) = \sum a_T \varsigma(T)T = 0$, so x is a linear span of trees T such that $\varsigma(T) = 0$. The converse is trivial. \square

We denote by $\mathcal{PT}_1^{(0)}(\mathcal{D})$ the set of partitioned trees $T \in \mathcal{PT}_1(\mathcal{D})$ with $\varsigma(T) = 0$. The preceding proposition implies that the Hopf algebras $(CP(\mathcal{D}), \cdot, \Delta)$ and $(\mathcal{H}_{CK}^{\mathcal{PT}_1^{(0)}(\mathcal{D})}, m, \Delta)$ are isomorphic. We obtain an explicit isomorphism between them:

Definition 33. Let $T \in \mathcal{PT}(\mathcal{D})$ and $\pi = \{P_1, \dots, P_k\}$ be a partition of $V(T)$. We shall write $\pi \triangleleft T$ if the following condition holds:

- For all $i \in [k]$, the partitioned rooted forest $T|_{P_i}$, denoted by T_i , belongs to $\mathcal{PT}_1^{(0)}(\mathcal{D})$.

If $\pi \triangleleft T$, the contracted graph T/π is a rooted forest (one forgets about the blocks of T). The vertex of T/π corresponding to P_i is decorated by T_i , making T/π an element of $\mathcal{T}(\mathcal{PT}_1^{(0)}(\mathcal{D}))$.

Corollary 34. The following map is a Hopf algebra isomorphism:

$$\Theta : \begin{cases} (CP(\mathcal{D}), \cdot, \Delta) & \longrightarrow (\mathcal{H}_{CK}^{\mathcal{PT}_1^{(0)}(\mathcal{D})}, \cdot, \Delta) \\ T \in \mathcal{PT}(\mathcal{D}) & \longrightarrow \sum_{\pi \triangleleft T} T/\pi. \end{cases}$$

Example 9. If $\mathcal{D} = \{1\}$, forgetting the decorations, with $a = \cdot$ and $b = \nabla$:

$$\Theta(\cdot) = \bullet_a, \quad \Theta(\dagger) = \dagger_a^a, \quad \Theta(\vee) = {}^a \vee_a^a, \quad \Theta(\nabla) = {}^a \vee_a^a + \bullet_b.$$

4.6 Extension of the preLie product \diamond to all partitioned trees

We now extend the preLie product \diamond to the whole $CP(\mathcal{D})$:

Proposition 35. *We define a product on $CP(\mathcal{D})$ in the following way:*

$$\forall T, T' \in \mathcal{PT}(\mathcal{D}), \quad T \diamond T' = \sum_{\substack{s \in V(T), \\ b \in \text{bl}(s) \sqcup \{*\}}} T \bullet_{s,b} T'.$$

Then $(CP(\mathcal{D}), \diamond, \cdot)$ is a Com-PreLie algebra.

Proof. Obviously, for any $x, y, z \in \mathcal{PT}(\mathcal{D})$, $(x \cdot y) \diamond z = (x \diamond z) \cdot x + x \cdot (y \diamond z)$. Let $T_1, T_2, T_3 \in \mathcal{PT}(\mathcal{D})$. Then:

$$\begin{aligned} (T_1 \diamond T_2) \diamond T_3 &= \sum_{\substack{s_1 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}}} \sum_{\substack{s_2 \in V(T_1), \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} (T_1 \bullet_{s_1, b_1} T_2) \bullet_{s_2, b_2} T_3 \\ &+ \sum_{\substack{s_1 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}}} \sum_{\substack{s_2 \in V(T_2), \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} (T_1 \bullet_{s_1, b_1} T_2) \bullet_{s_2, b_2} T_3 \\ &= \sum_{\substack{s_1 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}}} \sum_{\substack{s_2 \in V(T_1), \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} (T_1 \bullet_{s_1, b_1} T_2) \bullet_{s_2, b_2} T_3 \\ &+ \sum_{\substack{s_1 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}}} \sum_{\substack{s_2 \in V(T_2), \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} T_1 \bullet_{s_1, b_1} (T_2 \bullet_{s_2, b_2} T_3) \\ &= \sum_{\substack{s_1 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}}} \sum_{\substack{s_2 \in V(T_1), \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} (T_1 \bullet_{s_1, b_1} T_2) \bullet_{s_2, b_2} T_3 + T_1 \diamond (T_2 \diamond T_3). \end{aligned}$$

Hence:

$$\begin{aligned} (T_1 \diamond T_2) \diamond T_3 - T_1 \diamond (T_2 \diamond T_3) &= \sum_{\substack{s_1 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}}} \sum_{\substack{s_2 \in V(T_1), \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} (T_1 \bullet_{s_1, b_1} T_2) \bullet_{s_2, b_2} T_3 \\ &= \sum_{\substack{s_1 \neq s_2 \in V(T_1), \\ b_1 \in \text{bl}(s_1) \sqcup \{*\}, \\ b_2 \in \text{bl}(s_2) \sqcup \{*\}}} (T_1 \bullet_{s_1, b_1} T_2) \bullet_{s_2, b_2} T_3 \\ &+ \sum_{\substack{s \in V(T_1), \\ b_1 \neq b_2 \in \text{bl}(s) \sqcup \{*\}}} (T_1 \bullet_{s, b_1} T_2) \bullet_{s, b_2} T_3 + \sum_{\substack{s \in V(T_1), \\ b \in \text{bl}(s) \sqcup \{*\}}} (T_1 \bullet_{s, b} T_2) \bullet_{s, b} T_3. \end{aligned}$$

The three terms of this sum are symmetric in T_2, T_3 , so:

$$(T_1 \diamond T_2) \diamond T_3 - T_1 \diamond (T_2 \diamond T_3) = (T_1 \diamond T_3) \diamond T_2 - T_1 \diamond (T_3 \diamond T_2).$$

Finally, $(CP(\mathcal{D}), \diamond, \cdot)$ is Com-PreLie. □

Definition 36. Let $T = (t, I, d)$ and $T' = (t, I', d)$ be two elements of $\mathcal{PT}(\mathcal{D})$ with the same underlying decorated rooted trees. We shall say that $T \leq T'$ is I' is a refinement of I . This defines a partial order on $\mathcal{PT}(\mathcal{D})$.

Example 10. If $a, b, c, d \in \mathcal{D}$, $b \overset{c}{\underset{a}{\vee}} d \leq b \overset{c}{\underset{a}{\vee}} d, c \overset{b}{\underset{a}{\vee}} d, d \overset{b}{\underset{a}{\vee}} c \leq b \overset{c}{\underset{a}{\vee}} d$.

Theorem 37. *The following map is an isomorphism of Com-PreLie algebras:*

$$\Psi : \begin{cases} (CP(\mathcal{D}), \circ, \cdot) & \longrightarrow (CP(\mathcal{D}), \diamond, \cdot) \\ T \in \mathcal{PT}(\mathcal{D}) & \longrightarrow \sum_{T' \leq T} T'. \end{cases}$$

Proof. As \leq is a partial order, Ψ is bijective. Let $T_1, T_2 \in \mathcal{PT}(\mathcal{D})$.

1. If $T' \leq T_1 \cdot T_2$, let us put $T'_1 = T_1 \cap T'$ and $T'_2 = T_2 \cap T'$. Then, obviously, $T'_1 \leq T_1$ and $T'_2 \leq T_2$. Moreover, $T' = T'_1 \leq T'_2$. Conversely, if $T'_1 \leq T_1$ and $T'_2 \leq T_2$, then $T'_1 \cdot T'_2 \leq T_1 \cdot T_2$. Hence:

$$\Psi(T_1 \cdot T_2) = \sum_{T' \leq T_1 \cdot T_2} T' = \sum_{T'_1 \leq T_1, T'_2 \leq T_2} T'_1 \cdot T'_2 = \Psi(T_1) \cdot \Psi(T_2).$$

2. Let $s \in V(T_1)$ and $T' \leq T_1 \bullet_{s,*} T_2$. We put $T'_1 = T' \cap T_1$ and $T'_2 = T' \cap T_2$. Then, obviously, $T'_1 \leq T_1$ and $T'_2 \leq T_2$. If the block of roots of T_2 is also a block of T' , then $T' = T'_1 \bullet_{s,*} T'_2$. Otherwise, there exists a unique $b \in bl(s)$ such that $T' = T'_1 \bullet_{s,b} T'_2$. Conversely, if $T'_1 \leq T_1$, $T'_2 \leq T_2$, $s \in V(T'_1)$ and $b \in bl(s) \sqcup \{*\}$, then $T'_1 \bullet_{s,b} T'_2 \leq T_1 \bullet_{s,*} T_2$. Hence:

$$\begin{aligned} \Psi(T_1 \circ T_2) &= \sum_{s \in V(T_1)} \sum_{T' \leq T_1 \bullet_{s,*} T_2} T' \\ &= \sum_{T'_1 \leq T_1, T'_2 \leq T_2} \sum_{s \in V(T'_1), b \in bl(s) \sqcup \{*\}} T'_1 \bullet_{s,b} T'_2 \\ &= \Psi(T_1) \diamond \Psi(T_2). \end{aligned}$$

So Ψ is a Com-PreLie algebra isomorphism. □

Example 11. In the nondecorated case:

$$\begin{array}{ll} \Psi(\cdot) = \cdot, & \Psi(\dagger) = \dagger, \\ \Psi(\ddagger) = \ddagger, & \Psi(\blacktriangledown) = \blacktriangledown + 3\color{red}{\blacktriangledown} + \color{red}{\blacktriangledown}, \\ \Psi(\color{red}{\blacktriangledown}) = \color{red}{\blacktriangledown} + \color{red}{\blacktriangledown}, & \Psi(\color{red}{\blacktriangledown}) = \color{red}{\blacktriangledown} + \color{red}{\blacktriangledown}, \\ \Psi(\color{red}{\blacktriangledown}) = \color{red}{\blacktriangledown}, & \Psi(\color{red}{\blacktriangledown}) = \color{red}{\blacktriangledown}. \end{array}$$

References

- [1] Thomas Benes and Dietrich Burde, *Degenerations of pre-Lie algebras*, J. Math. Phys. **50** (2009), no. 11, 112102, 9.
- [2] D. J. Broadhurst and D. Kreimer, *Towards cohomology of renormalization: bigrading the combinatorial Hopf algebra of rooted trees*, Comm. Math. Phys. **215** (2000), no. 1, 217–236, arXiv:hep-th/0001202.
- [3] Alain Connes and Dirk Kreimer, *Hopf algebras, Renormalization and Noncommutative geometry*, Comm. Math. Phys **199** (1998), no. 1, 203–242.
- [4] ———, *Hopf algebras, renormalization and noncommutative geometry*, Comm. Math. Phys. **199** (1998), no. 1, 203–242.
- [5] Loïc Foissy, *The Hopf algebra of Fliess operators and its dual pre-Lie algebra*, Comm. Algebra **43** (2015), no. 10, 4528–4552.
- [6] ———, *A pre-Lie algebra associated to a linear endomorphism and related algebraic structures*, Eur. J. Math. **1** (2015), no. 1, 78–121.

- [7] Loïc Foissy, *Finite-dimensional comodules over the Hopf algebra of rooted trees*, J. Algebra **255** (2002), no. 1, 85–120.
- [8] W. Steven Gray and Luis A. Duffaut Espinosa, *A Faà di Bruno Hopf algebra for a group of Fliess operators with applications to feedback*, Systems Control Lett. **60** (2011), no. 7, 441–449.
- [9] Muriel Livernet, *A rigidity theorem for pre-Lie algebras*, J. Pure Appl. Algebra **207** (2006), no. 1, 1–18.
- [10] Jean-Louis Loday, *Scindement d’associativité et algèbres de Hopf*, Actes des Journées Mathématiques à la Mémoire de Jean Leray, Sémin. Congr., vol. 9, Soc. Math. France, Paris, 2004, pp. 155–172.
- [11] Jean-Louis Loday and Marí a Ronco, *On the structure of cofree Hopf algebras*, J. Reine Angew. Math. **592** (2006), 123–155.
- [12] Jean-Louis Loday and María Ronco, *Combinatorial Hopf algebras*, Quanta of maths, Clay Math. Proc., vol. 11, Amer. Math. Soc., Providence, RI, 2010, pp. 347–383.
- [13] J.-M. Oudom and D. Guin, *On the Lie enveloping algebra of a pre-Lie algebra*, J. K-Theory **2** (2008), no. 1, 147–167.
- [14] Jean-Michel Oudom and Daniel Guin, *Sur l’algèbre enveloppante d’une algèbre pré-Lie*, C. R. Math. Acad. Sci. Paris **340** (2005), no. 5, 331–336.